

AWARD NUMBER: W81XWH-14-1-0464

TITLE: The Roles of Primary Cilia in Cardiovascular System

PRINCIPAL INVESTIGATOR: Surya Nauli

CONTRACTING ORGANIZATION: Chapman University
Irvine, CA 92618-1908

REPORT DATE: October 2015

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE October 2015		2. REPORT TYPE Annual		3. DATES COVERED 15 Sep 2014 - 14 Sep 2015	
4. TITLE AND SUBTITLE The Roles of Primary Cilia in Cardiovascular System				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-14-1-0464	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Surya Nauli E-Mail: Nauli@chapman.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Chapman University 9401 Jeronimo Road Irvine, CA 92618-1908				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Hypertension and aneurysm are a prevalent problem in our society. Because the risk of these vascular diseases is too important to ignore, our studies are designed to provide exciting and original concepts to examine the etiologies of hypertension and aneurysm with regard to mechanosensory organelles (primary cilia). Especially within polycystic kidney disease, untreated hypertension can worsen kidney function. Aneurysm rupture in PKD patients also remains a devastating complication that often could result in stroke and death. Our studies therefore aim at investigating new ideas in understanding hypertension and aneurysm formation in PKD.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Unclassified	18. NUMBER OF PAGES 44	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
1. Introduction.....	3
2. Keywords.....	3
3. Accomplishments.....	3
4. Impact.....	6
5. Changes/Problems.....	6
6. Products.....	6
7. Participants & Other Collaborating Organizations.....	7
8. Special Reporting Requirements.....	7
9. Appendices.....	7

1. INTRODUCTION:

Polycystic kidney disease (PKD) is characterized by formation of fluid-filled cysts in both kidneys. PKD patients will eventually have renal failure, with subsequent dialysis or renal transplant. The genes mutated in PKD include Pkd1 and Pkd2, encoded for polycystin-1 and polycystin-2, respectively. Many studies have shown that the baseline circulating NO is much lower in PKD patients. This suggests possible vascular dysfunction. The purpose of our research is to investigate cilia function in the vascular endothelial cells. More important, hypertension has been a critically important risk factor for cardiovascular diseases in PKD patients, which occur early in these individuals, compared to their age-matched cohorts, and still remains the most frequent cause of mortality even when their renal function is still normal. With our expertise and tools available in our laboratory, we thus hypothesize that cilia in the cilia play crucial and important roles in regulating PKD pathology. Specifically, we will provide the first insights into physiological functions and cellular pathways of primary cilia in vasculatures and in PKD.

2. KEYWORDS:

Cardiovascular, cilia, ciliopathy, ciliotherapy, endothelia, epithelia, polycystic kidney disease, polycystin-1, polycystin-2, primary cilia

3. ACCOMPLISHMENTS:

Major Goals

The main goal of the project is to determine the roles of primary cilia in PKD. This goal is divided into two major subaims, and our Statement of Work (SOW) is as follow.

Aim 1 (months 1-30). We will study mechanosensory function of endothelial cilia in hypertension.

Aim 1.1 (months 1-12): We will measure blood pressure in cilium mutant mice in vivo.

Aim 1.2 (months 13-30): We will examine signaling mechanisms of cilia & their effects on blood pressure.

Aim 2 (months 7-36). We will study mechanosensory function of endothelial cilia in vascular aneurysm.

Aim 2.1 (months 7-20): We will quantify aneurysm formation in cilium mutant mice in vivo.



Aim 2.2 (months 20-36): We will identify signaling mechanisms of cilia & their consequence on aneurysm.

Accomplishments

Aim 1.1 (months 1-12): measurement of blood pressure

Over the past year, we have measured the systolic and diastolic blood pressure by non-invasive blood pressure system - tail cuff method with the aid of a computerized system (CODA system, Kent Scientific, Connecticut, USA). Measurements were performed at the baseline 3-times per week for 2 weeks after previous 3 days of training for each mouse. On each day of blood pressure measurement, 2 sets of 18 measurements were obtained including three measurements of training or acclimation. The measurements were averaged for each mouse with at least three mice for each genotype. All animals were tested by an investigator blinded to the genotypes of the animals. The data from the tail cuff method was also verified with limited studies with a more invasive, surgically implanted telemetry probe (data not shown).

When blood pressure was monitored in our mutant mice, *Survivin* knockout mice surprisingly did not show an elevated blood pressure (**Figure 1**). However, *Pkd1* and *Tg737* mice are hypertensive. Supporting this view, patients with PKD have a significantly greater chance to develop hypertension than general population.



Aim 2.1 (months 7-20): quantify aneurysm formation

We have started Aim 2.1 in the last couple of months. Our animal use for these studies has been approved by our Institutional Animal Care and Use Committee. We will use both traditional and conditional transgenic mouse models in our present studies. We have previously obtained traditional transgenic *Pkd*-mouse models *Pkd2^{+/-}* and *Tg737^{Orpk/Orpk}*. We will also use *survivin^{fllox/fllox}* and *Pkd1^{fllox/fllox}* conditional mouse models. To generate kidney-specific knockout, we have bred our mice with *Mx1Cre* mice. To generate vascular-specific knockout, we have used and confirmed *PdgfbCre* and *Tie2Cre* mice (**Figure 2**). Briefly, one-week old pups were injected intra-peritoneally with 62.5 µg of 50 µL polyinosinic:polycytidylic ribonucleic acid (pI:pC) every day for five consecutive days.

Tie2Cre or *PdgfbCre* mouse was bred with *Gt(ROSA)26Sor* (*Rosa26*) mouse, resulting in either *PdgfbCre•Rosa26* or *Tie2Cre•Rosa26* genotype. The *Rosa26* mouse was used as a control. These 3 mouse groups were induced to activate Cre which acts on *Rosa26* allele. The *Rosa26* genetic background is used as a

reporter system to verify and validate the efficiency of the *Cre* mice. The *Rosa26* includes its inducible fluorescence reporter system; i.e. all endothelia lining the vasculatures have red fluorescence (non-induced). The red fluorescence will be replaced with green fluorescence upon *Cre* recombinant (induction). Abdominal aortas were isolated, stained with nuclear marker (blue), and imaged for their green/red fluorescence (**Figure 2a**). Quantitation analysis of vascular-lining endothelia indicates high efficacy of both *PdgfbCre* and *Tie2Cre* backgrounds to delete a specific gene in vascular lining endothelia (**Figure 2b**). N=4 for each genotype and treatment.

Training and Professional Development

The project will continue to be part of our research-training program for two postdoctoral fellows and one graduate student. All of our trainees will continue working on the proposed studies, and results of their studies will be presented in conferences, such as FASEB meeting. This is an important meeting for our trainees to demonstrate their research productivity. We meet one on one in my office regularly to discuss any technical problems that may have arisen from their experiments. All trainees also participate in our laboratory meeting. All are expected to present their research progress in our laboratory meeting.

In addition, the Postdoctoral Research Committee and Graduate Committee will also evaluate the productivity of our postdoctoral fellows and graduate student, respectively. For most of the technical skills and methods, they will be trained within our laboratory and our surrounding laboratories. For more specialized skills, we will encourage them to attend various workshops. All trainees are recommended and student is required to the following courses.

- PHS601- Research Ethics and Regulations
- PHS 614- Biologics
- PHS 636- Proteomics

Results Dissemination

No outreach activity was attempted. Relevant results have been published and deposited into the National Library of Medicine.

1. Liu T, Jin X, Prasad RM, Sari Y, **Nauli SM**. Three types of ependymal cells with intracellular calcium oscillation are characterized by distinct cilia beating properties. *J Neurosci Res*. 2014 Sep;92(9):1199-204. [PMID:24811319]
2. Muntean BS, Jin X, Williams FE, **Nauli SM**. Primary cilium regulates CaV1.2 expression through Wnt signaling. *J Cell Physiol*. 2014 Dec;229(12):1926-34. [PMID:24700505]
3. Atkinson KF, Kathem SH, Jin X, Muntean BS, Abou-Alaiwi WA, Nauli AM, **Nauli SM**. Dopaminergic signaling within the primary cilia in the renovascular system. *Front Physiol*. 2015 Apr 16;6:103. [PMID:25932013]
4. Mohieldin AM, Zubayer HS, Al Omran AJ, Saternos HC, Zarban A, **Nauli SM**, AbouAlaiwi WA. Vascular Endothelial Primary Cilia: Mechanosensation and Hypertension. *Curr Hypertens Rev*. 2015 Jun 30 [In Press; PMID:26122329]

Future Goal

Our goal for next funding year is to complete Aim 2.1 and to initiate Aim 1.2.

4. IMPACT:

Impact on the discipline

Primary cilia are sensory organelles that extend from the cell surface and sense extracellular signals. Endothelial primary cilia protruding from the inner surface of blood vessel walls sense changes in blood flow and convert this mechanosensation into an intracellular biochemical/molecular signal, which triggers a cellular response. Endothelial cilia dysfunction may contribute to the impairment of this response and thus be directly implicated in the development of vascular abnormalities such as hypertension as seen in our PKD animal models. The completion of this project may present primary cilia as a novel therapeutic target for vascular hypertension.

Impact on other disciplines

Although our project is directed to lower blood pressure in patients with PKD, it may apply to a general population with hypertension. The completion of our project may serve as a proof of concept for targeted-clinical therapy on primary cilia as a novel mechanism in general hypertensive patients.

Impact on technology transfer

Nothing to Report at this time.

Impact on society

Nothing to Report.

5. CHANGES AND PROBLEMS:

Nothing to Report.

6. PRODUCTS:

1. Liu T, Jin X, Prasad RM, Sari Y, **Nauli SM**. Three types of ependymal cells with intracellular calcium oscillation are characterized by distinct cilia beating properties. *J Neurosci Res*. 2014 Sep;92(9):1199-204. [PMID:24811319]
2. Muntean BS, Jin X, Williams FE, **Nauli SM**. Primary cilium regulates CaV1.2 expression through Wnt signaling. *J Cell Physiol*. 2014 Dec;229(12):1926-34. [PMID:24700505]
3. Atkinson KF, Kathem SH, Jin X, Muntean BS, Abou-Alaiwi WA, Nauli AM, **Nauli SM**. Dopaminergic signaling within the primary cilia in the renovascular system. *Front Physiol*. 2015 Apr 16;6:103. [PMID:25932013]
4. Mohieldin AM, Zubayer HS, Al Omran AJ, Saternos HC, Zarban A, **Nauli SM**, AbouAlaiwi WA. Vascular Endothelial Primary Cilia: Mechanosensation and Hypertension. *Curr Hypertens Rev*. 2015 Jun 30 [In Press; PMID:26122329]

7. PARTICIPANTS:

Individuals

Name:	<i>Kimberly Atkinson</i>
Project Role:	<i>Postdoctoral Fellow</i>
Researcher Identifier (e.g. ORCID ID):	<i>N/A</i>
Nearest person month worked:	<i>~9</i>
Contribution to Project:	<i>Dr. Atkinson has been working on the in vivo mouse model.</i>
Funding Support:	<i>This award</i>

Name:	<i>Pala Rajasekharreddy</i>
Project Role:	<i>Postdoctoral Fellow</i>
Researcher Identifier (e.g. ORCID ID):	<i>N/A</i>
Nearest person month worked:	<i>~6</i>
Contribution to Project:	<i>Dr. Rajasekharreddy has been working on the in vitro assay.</i>
Funding Support:	<i>This award</i>

Name:	<i>Rinzhin Sherpa</i>
Project Role:	<i>Graduate Student</i>
Researcher Identifier (e.g. ORCID ID):	<i>N/A</i>
Nearest person month worked:	<i>~9</i>
Contribution to Project:	<i>Mr. Sherpa has been working on the in vivo mouse model and in vitro cilia measurement.</i>
Funding Support:	<i>This award</i>

Change in Other Supports or Key Personnel

Nothing to Report

Partner Organization

Nothing to Report

8. SPECIAL REPORTING REQUIREMENTS:

Not applicable

9. APPENDICES:

See Below

Three Types of Ependymal Cells With Intracellular Calcium Oscillation Are Characterized by Distinct Cilia Beating Properties

Tongyu Liu, Xingjian Jin, Rahul M. Prasad, Youssef Sari, and Surya M. Nauli*

Department of Pharmacology, University of Toledo, Toledo, Ohio

Ependymal cells are multiciliated epithelial cells that line the ventricles in the adult brain. Abnormal function or structure of ependymal cilia has been associated with various neurological deficits. For the first time, we report three distinct ependymal cell types, I, II, and III, based on their unique ciliary beating frequency and beating angle. These ependymal cells have specific localizations within the third ventricle of the mouse brain. Furthermore, neither ependymal cell types nor their localizations are altered by aging. Our high-speed fluorescence imaging analysis reveals that these ependymal cells have an intracellular pacing calcium oscillation property. Our study further shows that alcohol can significantly repress the amplitude of calcium oscillation and the frequency of ciliary beating, resulting in an overall decrease in volume replacement by the cilia. Furthermore, the pharmacological agent cilostazol could differentially increase cilia beating frequency in type II, but not in type I or type III, ependymal cells. In summary, we provide the first evidence of three distinct types of ependymal cells with calcium oscillation properties. © 2014 Wiley Periodicals, Inc.

Key words: cilia; cerebrospinal fluid; calcium

Cilia are generally classified as solitary nonmotile and bundled motile organelles (Abou Alaiwi et al., 2009a; Nauli et al., 2011). Motile and nonmotile cilia have been implicated in fundamental processes of development and disease. Motile cilia can be found in the ependymal cells, forming a lining in the brain ventricles and central canal of the spinal cord. Ependymal cells are ciliated, simple cuboidal, epithelium-like glial cells that move cerebrospinal fluid (CSF) along the ventricles (Del Bigio, 1995). Abnormal ependymal cilia result in hydrocephalus induced by anomalous flow of CSF (Banizs et al., 2005; Baas et al., 2006; Wodarczyk et al., 2009; Tissir et al., 2010).

Ependymal cells also play an important role in regulating pluripotent neural stem cells (Rietze et al., 2001). Beating of ependymal cilia is required for normal CSF flow, which functions as a guide for specific directional migration of new neurons (Sawamoto et al., 2006). The coupling between ependymal cilia beating and hydrodynamic forces has been proposed to regulate planar cell

polarity during development or stroke (Guirao et al., 2010; Mirzadeh et al., 2010; Devaraju et al., 2013). In addition, ependymal cilia play major roles in CSF dynamics, cerebral fluid balance, secretion, toxin metabolism, and many other functions (Genzen et al., 2009; Appelbe et al., 2013). Although ependymal cells regulate CSF flow, which regulates many neuronal processes, different types of ependymal cells have not been distinguished. We show here, based on the cilia beating frequency and beating angle, that ependymal cells can be distinctly categorized into three types. Furthermore, each type of ependymal cell is uniquely localized within the ventricle.

MATERIALS AND METHODS

All animal experiments were approved by the University of Toledo's Institutional Animal Care and Use Committee (IACUC). The wild-type mice were euthanized with carbon dioxide for 5 min. After craniotomy, the whole brain was removed. The sagittal slice was dissected with a thickness of about 100 μ m and was immediately embedded in Dulbecco's modified Eagle's medium (DMEM; Cellgro Corning Life Sciences-Mediatech, Manassas, VA) at 39°C in the presence of 95%/5% O₂/CO₂ mixture.

Immunofluorescence Microscopy

The brain slice was fixed in phosphate buffer containing 3% paraformaldehyde and 2% sucrose for 10 min. Mouse primary antibody antiacetylated α -tubulin was used at a dilution of 1:5,000 (Sigma, St. Louis, MO). The brain slice was

Additional Supporting Information may be found in the online version of this article.

Contract grant sponsor: NIH, contract grant number: DK080640; Contract grant sponsor: DOD, contract grant number: PR130153; Contract grant sponsor: University of Toledo.

*Correspondence to: Surya M. Nauli, PhD, Department of Pharmacology, University of Toledo, MS 1015, Health Education Building, Room 274, 3000 Arlington Ave., Toledo, OH 43614.

E-mail: surya.nauli@utoledo.edu

Received 15 January 2014; Revised 7 April 2014; Accepted 7 April 2014

Published online 9 May 2014 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/jnr.23405

incubated in primary antibody solution overnight. Secondary antibody fluorescein anti-mouse IgG was used at a dilution of 1:500 (Vector Laboratories, Burlingame, CA), and the brain slice was incubated with secondary antibody solution for 1 hr. Before observation under a TiU fluorescent microscopy (Nikon, Tokyo, Japan), the section was dyed with DAPI for 5 min (Abou Alaiwi et al., 2014).

Measurement of Cilia Beating Frequency

The prepared brain slice was kept in a customized glass-bottom plate covered with 500 μ l DMEM containing 2% B27 at 39°C. In some cases, 0.25% ethanol was added to the media. The video of cilia beating was captured with a TiU high-resolution differentiation interference contrast microscope. The capture rate of the video was 5 msec for a minimum of 1 sec (200 frames per sec).

Measurement of Fluid Movement and Volume Replacement

Because of the transparency of the buffer solution, we used 200-nm latex beads (Molecular Probes, Eugene, OR) to help analyze speed in the solution movement. The velocity of fluid movement was calculated by tracing one single nanobead flowing across the third ventricle wall. The overall fluid volume moved by ependymal cilia was calculated with the following formula: volume replacement ($\mu\text{m}^3/\text{stroke}$) = fluid movement velocity ($\mu\text{m}^3/\text{sec}$)/cilia beating frequency (stroke/sec).

Calcium Signal Recording

To record cytosolic calcium oscillation, the brain slice was incubated with 20 $\mu\text{g}/\text{ml}$ fluo-2 (TEFLabs, Austin, TX) for 30 min at 39°C. The tissue was then transferred to a glass-bottom plate covered with 500 μ l DMEM containing 2% B27 (Gibco, Rockville, MD) at 39°C. In some cases, 0.25% ethanol was added to the media. The video of calcium oscillation was recorded at a capture rate of 5 msec for a minimum of 1 sec (200 frames per sec), with excitation and emission wavelengths of 488 nm and 515 nm, respectively.

RESULTS

Ependymal Cells Can Be Classified Into Three Types Based on Their Cilia Beating Frequency

We cut the mouse brain in a sagittal plane to enhance our observation of the entire third ventricle. To verify our high-resolution differential interference contrast and fluorescence microscope systems, we examined the presence of ependymal cilia in the third ventricle (Fig. 1). Ependymal cilia were confirmed with a ciliary marker, acetylated- α -tubulin. Although in the control permeabilized brain no fluid movement was observed in nonbeating ependymal cilia (Supp. Info. Movie 1), we could observe the direction of fluid movement via oil ink in a freshly prepared brain ex vivo (Supp. Info. Movie 2).

We next attempted to quantify the cilia beating frequency. Based on our observation of individual ependymal cells from 87 independent experiments, we were surprised to notice that there were wide variations in the beating fre-




Fig. 1. The presence of ependymal cilia in mouse brain. Shown here are ependymal cells from the third ventricle of a mouse brain. The brain section was stained with anti-acetylated- α -tubulin, a ciliary marker (green), and counterstained with DAPI, a nucleus marker (blue). Individual differential interference contrast (**top**), fluorescence (**middle**), and merged (**bottom**) images are shown. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]




Fig. 2. Categories of ependymal cells in mouse brain. Based on a total of 87 independent experiments or preparations (represented as dots), ependymal cells could be classified into three types (grouped within circle or oval lines). Type I ependymal cilia had the highest beating frequency (>60 Hz), with a ciliary beating angle of less than 90° . Type II ependymal cilia had a medium beating frequency (30–60 Hz), with a ciliary beating angle between 90° and 135° . Type III ependymal cilia had the slowest beating frequency (<30 Hz), with a ciliary beating angle of greater than 135° . The cartoon on the right depicts the beating angles of ependymal cilia.

quencies of ependymal cilia (Fig. 2). We could accurately assign each ependymal cell to one of three classifications depending on its ciliary beating. Type I ependymal cells




Fig. 3. Differential ependymal cell types and localization. Based on the ependymal cilia beating frequency, different ependymal cell types were enriched at certain locations within the third ventricle. A freshly cut brain section was imaged to show the third ventricle region (top). The cartoon (middle) and brain section (bottom) of the third ventricle depict localizations of different ependymal cells from a sagittal view. Scale bar = 30 μ m. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

had the highest beating frequency (>60 Hz) and had a beating angle of less than 90° (Supp. Info. Movie 3). Type II ependymal cells had a medium beating frequency (30–60 Hz) and had a beating angle between 90° and 135° (Supp. Info. Movie 4). Type III ependymal cells had the slowest beating frequency (<30 Hz) and had a beating angle of more than 135° (Supp. Info. Movie 5).

To understand the difference in localizations among these ependymal cells, we mapped the distributions of ependymal cell types within the third ventricle (Fig. 3). Our mapping analysis indicated that type I cells were widely distributed along the ventricle walls, but they were absent at both inferior and superior corners of the third ventricle. Type II cells were observed mainly on the anterior wall of the ventricle, but they could also be found at the posterior wall of the ventricle. Type III cells were distributed almost exclusively in the inferior and superior corners of the third ventricle. Of note is that we could find all three cell types at the lower wall of the third ventricle.

Cilia Beating Frequency Is Age Independent in Third Ventricles

Because ependymal cilia have been proposed to move CSF to support migration of various substances




Fig. 4. Ependymal cilia beating frequency in different mouse age groups. All types of ependymal cilia in control and alcohol-treated mouse third ventricles were studied at different ages. Although beat frequency of ependymal cilia was not affected by age, acute alcohol treatment significantly decreased cilia beating frequency in all types of ependymal cells. At least three independent preparations were used for each ependymal cell type and age group.

(Banizs et al., 2005; Sawamoto et al., 2006), we next sought to understand the role of aging in ependymal cilia beating. Mice were grouped according to age: 3–5 weeks, 9–11 weeks, and 15–17 weeks. The data indicate that we could distinctively classify the ependymal cilia beating into three types regardless of the age groups (Fig. 4). More importantly, there was no evidence that age was a factor in regulating ependymal cilia beating.

Ependymal Cilia Beating Can Be Repressed by Ethanol

To confirm further that our classification of ependymal cells was valid and consistent, we performed chemical screening on the cilia beating frequency. Ethanol at a concentration of 0.25% provided us with the most consistent changes in ependymal cilia beating. The data indicated that ethanol repressed ependymal cilia beating regardless of the age group (Fig. 4). Most importantly, ethanol repressed




Fig. 5. Effects of alcohol on the dynamics of mouse third ventricle. The ex vivo brain slice was incubated without (control) or with (ethanol) alcohol for 5 min. **a:** Compared with control, alcohol treatment significantly decreased cilia beating frequency. **b:** This resulted in a decrease in fluid movement, as indicated by speed of fluid movement surrounding the ependymal cilia. **c:** Further calculation of the volume replacement and cilia beating indicated that, compared with control, alcohol significantly decreased volume replacement for each stroke of cilia beating. This indicates that alcohol not only decreases the frequency of ependymal cilia beating but also reduces the efficiency of each cilia stroke. At least 10 independent preparations were used for each ependymal cell type and treatment group.

ependymal cilia beating (Fig. 5a), resulting in a significant decrease in fluid movement velocity around ependymal cilia (Fig. 5b). Because of the transparency of the fluid media, we used nanobeads to guide us in measuring the speed of the fluid movement (Supp. Info. Movie 6). Given the fluid movement velocity, we estimated the volume replacement for each stroke of cilia beating efficiency. Our calculation indicated that ethanol not only repressed cilia beating frequency but also decreased the efficiency of ependymal cilia to move fluid per each stroke (Fig. 5c).

Calcium Signaling by Ependymal Cilia Can Be Altered by Alcohol

Fluid-shear stress resulting from fluid movement above a layer of cells can generate intracellular calcium

signaling (Abou Alaiwi et al., 2009b; Jin et al., 2013). We therefore examined the calcium signaling within the third ventricle (Fig. 6a). As expected, we did not see any apparent calcium oscillation in fixed brain sections (Supp. Info. Movie 7). In the absence (Supp. Info. Movie 8) or presence (Supp. Info. Movie 9) of ethanol, however, calcium oscillation was observed on ependymal cells. The frequency of calcium oscillation was not changed either by mock (PBS or control) or by ethanol treatment (Fig. 6b). Although the frequency of calcium signal in each ependymal cell type was unchanged, the amplitude of calcium signal was significantly repressed in the ethanol group compared with the control groups of each of the corresponding ependymal cell types (Fig. 6c).

DISCUSSION

We report here for the very first time that there are three distinct types of ependymal cells uniquely and specifically positioned within the third ventricle. We classified them based on their cilia beating frequency as type I (>60Hz), type II (30–60 Hz), and type III (<30 Hz). The beat frequency for each type of ependymal cilia is age independent. We also report here that ependymal cells are characterized by calcium oscillations, the frequency and amplitude of which are the same in all ependymal cell types. Our chemical testing indicates that alcohol has a profound effect on the beating frequency of the ependymal cilia, resulting in a significant decrease in fluid movement and volume replacement. Although alcohol did not change the frequency of calcium oscillation in the ependymal cells, the amplitude of calcium oscillation was significantly repressed.

Even with the advancement in the technology of high-speed digital imaging (Lehtreck et al., 2009), there has been no report on different types of ependymal cells. We thus believe that our study is the first to identify distinct ependymal cilia, which is fundamentally important to gain basic understanding of ependymal physiology. For example, many substances that are known to alter cilia beating (Sisson et al., 1991; Sisson, 1995) may fail to show an effect in ependymal cells, especially when ependymal cells are randomly analyzed (Smith et al., 2013). To validate our point further, 1% ethanol was reported to have no effect on the beating frequency of ependymal cilia (Smith et al., 2013). After we classified the ependymal cilia into three types, however, our data clearly showed that ethanol as low as 0.25% had a definitive effect on ependymal cilia beating frequency.

Our study also revealed a unique aspect of calcium signaling in ependymal cells. It was previously thought that cardiac myocytes were the only cells that naturally have a pacing calcium oscillation. We used a fluorescence high-speed digital imaging system to demonstrate that, like myocytes, ependymal cells also have an oscillating intracellular calcium pattern. The frequency and amplitude of this calcium oscillation are similar among all three types of ependymal cells, indicating that the calcium may reflex functional states rather than types of the ependymal




Fig. 6. Effects of alcohol on calcium oscillation in mouse brain ependymal cells. After being loaded with calcium indicator fluo-2, the ex vivo brain slice was incubated without (control) or with (ethanol) alcohol for 5 min. **a:** Intracellular calcium of ependymal cells was measured every 5 msec, as indicated by the representative blue and red lines. **b:** There was no difference in calcium oscillation frequency

between control and alcohol-treated groups. **c:** However, the amplitude of calcium oscillation was significantly repressed in alcohol-treated groups compared with control groups in all types of ependymal cells. At least five independent preparations were used for each ependymal cell type and treatment group. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

cells. Furthermore, the frequencies of ciliary beating and calcium oscillation on ependymal cells might not be associated with one another. Consistent with this idea, an ethanol-induced decrease in ciliary beating frequency did not alter the frequency of calcium oscillation.

Despite the fact that the characteristics of cilia beating are very different among these three types of ependymal cilia, it is important to note that our understanding of these cells is just starting to unfold. For example, cilostazol could differentially increase cilia beating frequency in type II but not in type I or type III cells (Supp. Info. Fig. 1). Cilostazol is a specific inhibitor for phosphodiesterase-3, an enzyme that metabolizes cAMP to AMP, and it also regulates intracellular calcium (Kawanabe et al., 2012). It has been known that cAMP and calcium could regulate cilia beating frequency (Nguyen et al., 2001; Monkkonen et al., 2008; Genzen et al., 2009), but the differential effect of cilostazol and many other pharmacological agents requires further in-depth study to advance our understanding of the molecular and cellular biology of different ependymal cell types.

It is also worth mentioning that, among the three types of ependymal cells, type III cells were the most efficient at moving fluid volume with each ciliary stroke. Although their frequency of ciliary beating was the slowest, type III cells had the largest angle of stroke. The angle of stroke might therefore contribute significantly to mov-

ing fluid volume. However, it is important to mention that ciliary beat frequency was also a critical contributing factor in moving fluid volume, as seen with the ethanol treatment. Treatment with ethanol decreased fluid replacement significantly, primarily because of a slowing of ciliary beating.

Alcohol can produce a variety of detrimental effects in the central nervous system, leading to a wide range of impairments. Within minutes of alcohol consumption, the alcohol in the CSF reaches the same level as that in the blood (Kiianmaa and Virtanen, 1978; Agapejev et al., 1992; Huang and Huang, 2007). However, the effect of alcohol on each type of ependymal cell had never before been examined, although abnormal ependymal cilia are associated with ventricle enlargement associated with hydrocephalus (Banizs et al., 2005; Baas et al., 2006; Wodarczyk et al., 2009; Tissir et al., 2010). Consistent with this notion, the brains of alcoholics are known to have an increase in the size of the ventricles, causing hydrocephalus ex vacuo (de la Monte, 1988). The use of 0.25% of ethanol in our study was within the range of alcohol levels in the CSF as observed in humans and various animal models (Kiianmaa and Virtanen, 1978; Agapejev et al., 1992; Huang and Huang, 2007). Thus, our study also reflects a serious clinical implication in alcohol abusive behavior with regard to ventricle-lining ependymal cells, in addition to providing fundamental basic

scientific understanding of ependymal cilia and calcium signaling.

ACKNOWLEDGMENTS

We thank Charisse Montgomery for her editing service. T. Liu's work partially fulfilled the requirements for a Master's degree in Pharmacology.

REFERENCES

- Abou Alaiwi WA, Lo ST, Nauli SM. 2009a. Primary cilia: highly sophisticated biological sensors. *Sensors* 9:7003–7020.
- Abou Alaiwi WA, Takahashi M, Mell BR, Jones TJ, Ratnam S, Kolb RJ, Nauli SM. 2009b. Ciliary polycystin-2 is a mechanosensitive calcium channel involved in nitric oxide signaling cascades. *Circ Res* 104:860–869.
- Abou Alaiwi WA, Muntean BS, Ratnam S, Joe B, Liu L, Booth RL, Rodriguez I, Herbert BS, Bacallao RL, Fruttiger M, Mak TW, Zhou J, Nauli SM. 2014. Survivin-induced abnormal ploidy contributes to cystic kidney and aneurysm formation. *Circulation* 126:660–675.
- Agapejev S, Vassiliev I, Curi PR. 1992. Alcohol levels in cerebrospinal fluid and blood samples from patients under pathological conditions. *Acta Neurol Scand* 86:496–500.
- Appelbe OK, Bollman B, Attarwala A, Tribes LA, Muniz-Talavera H, Curry DJ, Schmidt JV. 2013. Disruption of the mouse *Jhy* gene causes abnormal ciliary microtubule patterning and juvenile hydrocephalus. *Dev Biol* 382:172–185.
- Baas D, Meinie A, Benadiba C, Bonnafé E, Meinie O, Reith W, Durand B. 2006. A deficiency in RFX3 causes hydrocephalus associated with abnormal differentiation of ependymal cells. *Eur J Neurosci* 24:1020–1030.
- Banizs B, Pike MM, Millican CL, Ferguson WB, Komlosi P, Sheetz J, Bell PD, Schwiebert EM, Yoder BK. 2005. Dysfunctional cilia lead to altered ependyma and choroid plexus function, and result in the formation of hydrocephalus. *Development* 132:5329–5339.
- de la Monte SM. 1988. Disproportionate atrophy of cerebral white matter in chronic alcoholics. *Arch Neurol* 45:990–992.
- Del Bigio MR. 1995. The ependyma: a protective barrier between brain and cerebrospinal fluid. *Glia* 14:1–13.
- Devaraju K, Barnabe-Heider F, Kokaia Z, Lindvall O. 2013. FoxJ1-expressing cells contribute to neurogenesis in forebrain of adult rats: evidence from in vivo electroporation combined with piggyBac transposon. *Exp Cell Res* 319:2790–2800.
- Genzen JR, Yang D, Ravid K, Bordey A. 2009. Activation of adenosine A2B receptors enhances ciliary beat frequency in mouse lateral ventricle ependymal cells. *Cerebrospinal Fluid Res* 6:15.
- Guirao B, Meunier A, Mortaud S, Aguilar A, Corsi JM, Strehl L, Hirota Y, Desoeuvre A, Boutin C, Han YG, Mirzadeh Z, Cremer H, Montcouquiol M, Sawamoto K, Spassky N. 2010. Coupling between hydrodynamic forces and planar cell polarity orients mammalian motile cilia. *Nat Cell Biol* 12:341–350.
- Huang CM, Huang RH. 2007. Ethanol inhibits the sensory responses of cerebellar granule cells in anesthetized cats. *Alcoholism Clin Exp Res* 31:336–344.
- Jin X, Mohieldin AM, Muntean BS, Green JA, Shah JV, Mykityn K, Nauli SM. 2013. Cilioplasm is a cellular compartment for calcium signaling in response to mechanical and chemical stimuli. *Cell Mol Life Sci* doi: 10.1007/s00018-013-1483-1.
- Kawanabe Y, Takahashi M, Jin X, Abdul-Majeed S, Nauli AM, Sari Y, Nauli SM. 2012. Cilostazol prevents endothelin-induced smooth muscle constriction and proliferation. *PLoS One* 7:e44476.
- Kiianmaa K, Virtanen P. 1978. Ethanol and acetaldehyde levels in cerebrospinal fluid during ethanol oxidation in the rat. *Neurosci Lett* 10:181–186.
- Lehtreck KF, Sanderson MJ, Witman GB. 2009. High-speed digital imaging of ependymal cilia in the murine brain. *Methods Cell Biol* 91:255–264.
- Mirzadeh Z, Han YG, Soriano-Navarro M, Garcia-Verdugo JM, Alvarez-Buylla A. 2010. Cilia organize ependymal planar polarity. *J Neurosci* 30:2600–2610.
- Monkkonen KS, Hirst RA, Laitinen JT, O'Callaghan C. 2008. PACAP27 regulates ciliary function in primary cultures of rat brain ependymal cells. *Neuropeptides* 42:633–640.
- Nauli SM, Haymour HS, AbouAlaiwi WA, Lo ST, Nauli AM. 2011. Primary cilia are mechanosensory organelles in vestibular tissues. mechanosensitivity and mechanotransduction. In: Kamkin A, Kiseleva I, editors. *Mechanosensitivity and mechanotransduction*. New York: Springer. Chapter 14.
- Nguyen T, Chin WC, O'Brien JA, Verdugo P, Berger AJ. 2001. Intracellular pathways regulating ciliary beating of rat brain ependymal cells. *J Physiol* 531:131–140.
- Rietze RL, Valcanis H, Brooker GF, Thomas T, Voss AK, Bartlett PF. 2001. Purification of a pluripotent neural stem cell from the adult mouse brain. *Nature* 412:736–739.
- Sawamoto K, Wichterle H, Gonzalez-Perez O, Cholfin JA, Yamada M, Spassky N, Murcia NS, Garcia-Verdugo JM, Marin O, Rubenstein JL, Tessier-Lavigne M, Okano H, Alvarez-Buylla A. 2006. New neurons follow the flow of cerebrospinal fluid in the adult brain. *Science* 311:629–632.
- Sisson JH. 1995. Ethanol stimulates apparent nitric oxide-dependent ciliary beat frequency in bovine airway epithelial cells. *Am J Physiol* 268:L596–L600.
- Sisson JH, Tuma DJ, Rennard SI. 1991. Acetaldehyde-mediated cilia dysfunction in bovine bronchial epithelial cells. *Am J Physiol* 260:L29–L36.
- Smith CM, Radhakrishnan P, Sikand K, O'Callaghan C. 2013. The effect of ethanol and acetaldehyde on brain ependymal and respiratory ciliary beat frequency. *Cilia* 2:5.
- Tissir F, Qu Y, Montcouquiol M, Zhou L, Komatsu K, Shi D, Fujimori T, Labeau J, Tyteca D, Courtoy P, Poumay Y, Uemura T, Goffinet AM. 2010. Lack of cadherins *Celsr2* and *Celsr3* impairs ependymal ciliogenesis, leading to fatal hydrocephalus. *Nat Neurosci* 13:700–707.
- Wodarczyk C, Rowe I, Chiaravalli M, Pema M, Qian F, Boletta A. 2009. A novel mouse model reveals that polycystin-1 deficiency in ependyma and choroid plexus results in dysfunctional cilia and hydrocephalus. *PLoS One* 4:e7137.

Primary Cilium Regulates CaV1.2 Expression Through Wnt Signaling

BRIAN S. MUNTEAN,¹ XINGJIAN JIN,² FREDERICK E. WILLIAMS,² AND SURYA M. NAULI^{1,2*}

¹Department of Medicinal and Biological Chemistry, The University of Toledo, Toledo, Ohio

²Department of Pharmacology, The University of Toledo, Toledo, Ohio

Primary cilia are sensory organelles that provide a feedback mechanism to restrict Wnt signaling in the absence of endogenous Wnt activators. Abnormal Wnt signaling has been shown to result in polycystic kidney disease (PKD) although the exact mechanism has been debated. Previously, we reported that the calcium channel CaV1.2 functions in primary cilia. In this study, we show that CaV1.2 expression level is regulated by Wnt signaling. This occurs through modulation of mitochondrial mass and activity resulting in increased reactive oxygen species which generate oxidative DNA lesions. We found that the subsequent cellular DNA damage response triggers increased CaV1.2 expression. In the absence of primary cilia where Wnt signaling is upregulated, we found that CaV1.2 is overexpressed as a compensatory mechanism. We show for the first time that CaV1.2 knockdown in zebrafish results in classic primary cilia defects including renal cyst formation, hydrocephalus, and left-right asymmetry defects. Our study shows that suppressed Wnt signaling prevents CaV1.2 expression ultimately resulting in PKD phenotypes. Thus, CaV1.2 expression is tightly regulated through Wnt signaling and plays an essential sensory role in primary cilia necessary for cellular homeostasis.

J. Cell. Physiol. 229: 1926–1934, 2014. © 2014 Wiley Periodicals, Inc.

Wnt signaling is an important regulator of cellular development and proliferation. In the absence of Wnt ligands, a complex consisting of Axin, adenomatous polyposis coli (APC), and glycogen synthase kinase 3 β (GSK3 β) induces β -catenin for ubiquitylation by SCF E3 ligases and eventual proteasomal degradation (Aberle et al., 1997). Wnt signal transduction occurs when secreted Wnt ligands bind Frizzled receptors resulting in phosphorylation of LRP5/6. The Axin-APC-GSK3 β complex is then recruited to LRP5/6 at the cell membrane which prevents β -catenin from being degraded. The accumulated β -catenin translocates to the nucleus and activates transcription of Wnt target genes (Muntean et al., 2012).

Primary cilia are non-motile sensory organelles present as a single copy on most differentiated cells in the body. Calcium signaling through primary cilium is essential for renal epithelial homeostasis (Nauli et al., 2003; Jin et al., 2013). Cilia extend from the cell surface through the basal body via intraflagellar transport (Moyer et al., 1994). The most common pathologies resulting from cilia dysfunction include polycystic kidney (Wilson, 2004), hypertension (Nauli et al., 2008; AbouAlaiwi et al., 2009), aneurysm (AbouAlaiwi et al., 2013), hydrocephalus (Carter et al., 2012), and left-right asymmetry defects (Norris, 2012).

Abnormal Wnt signaling has also been linked to polycystic kidney disease (PKD) (Lancaster et al., 2009). For example, increased cytosolic and nuclear β -catenin accumulation has been shown in various cilia mutant cells (Gerdes et al., 2007; Lancaster et al., 2011). Thus, primary cilia are thought to provide a feedback mechanism that restricts Wnt signaling in the absence of appropriate ligands (Gerdes et al., 2007; Lancaster et al., 2009, 2011).

We recently showed that voltage-gated L-type calcium channel CaV1.2 localized to primary cilia in renal epithelia (Jin et al., 2013). Because Wnt signaling has also been reported to modulate mitochondrial physiology (Yoon et al., 2010), we hypothesized that primary cilia play a role in Wnt regulation of mitochondria through CaV1.2. We show that although CaV1.2 is not required for cilia formation, Wnt increases mitochondria mass and activity in CaV1.2 deficient renal epithelial cells. This increases mitochondria reactive oxidative species (ROS) and DNA damage, resulting in PKD phenotypes. Thus, our study suggests that primary cilia may play a role in CaV1.2 expression level through Wnt regulation of mitochondria.

Materials and Methods

The experimental use of zebrafish was approved by The University of Toledo's Institutional Animal Care and Use Committee (IACUC). The use of lentiviral components was approved by the Institutional Biosafety Committee of The University of Toledo.

Cell culture

Immortalized mouse renal epithelial wild-type and *Tg737^{orp/k}* cells were cultured in Dulbecco's Modified Eagle Medium (Corning Cellgro) supplemented with 10% fetal bovine serum (HyClone Laboratories, Logan, Utah) and 1% penicillin/streptomycin (Corning Cellgro) at 39°C in 5% CO₂, as previously described (AbouAlaiwi et al., 2013). Prior to experiments, cells were treated with 100 ng/ml recombinant Wnt3a (R&D Systems, Minneapolis, MN) for 3 days and serum starved for 24 h.

RNAi knockdown cells

shRNA lentiviral vectors specific to *Cacna1c* (Origene; pGFP-C-shLenti clone ID: TL500242) were transfected into HEK293T cells. Viral supernatants were collected after 48 h, centrifuged, and passed through a 0.45 μ m filter. Cells were then spin-inoculated with pseudoviral particles containing 8 μ g/ml polybrene at 2,500 rpm for 30 min at 30°C and then cultured for up to 1 week. CaV1.2 knockdown was verified through Western blot analysis.

Brian S. Muntean and Xingjian Jin contributed equally to this work.

* Correspondence to: Surya M. Nauli, Department of Pharmacology, The University of Toledo, MS 1015, Health Education Building; Room 274, 3000 Arlington Ave, Toledo, OH 43614.

E-mail: surya.nauli@utoledo.edu

Manuscript Received: 20 January 2014

Manuscript Accepted: 1 April 2014

Accepted manuscript online in Wiley Online Library (wileyonlinelibrary.com): 2 April 2014.

DOI: 10.1002/jcp.24642

TABLE 1. shRNA sequences

Descriptions	Sequences
Scrambled control	5'-TGACCACCCTGACCTACGGCGTGCAGTGC-3'
<i>Cacna1c</i>	5'-TCAGAGTGCCTCACTGTTCTCGTGACCT-3'

Stable knockdown cell lines were obtained through puromycin selection. The following shRNA sequences were used (Table 1).

Immunostaining studies

Cells were grown to confluence on collagen-coated glass coverslips and differentiated in serum-free media for 24 h. Cells were then fixed in 4% paraformaldehyde in PBS containing 2% sucrose, permeabilized in 10% triton X-100, incubated sequentially with primary followed by secondary antibodies for 1 h each, and finally mounted on a slide with DAPI hard set mounting media (Vector Laboratories, Burlingame, CA). The following primary antibody dilutions were used: acetylated- α -tubulin 1:10,000 (Sigma-Aldrich, St. Louis, MO) and CaV1.2 1:50 (Alomone Labs, Jerusalem, Israel). Anti-mouse Texas Red and anti-rabbit FITC fluorescent conjugated secondary antibodies were used at 1:500 (VectorLabs).

Mitochondrial studies

MitoTracker Green FM and MitoTracker Red CMXRos (Cell Signaling Technology) were incubated with cells at 100 nM for 30 min at 37°C. MitoSOX (Life Technologies) was incubated with cells at 5 μ M for 10 min at 37°C. After staining, cells were washed three times with PBS and analyzed immediately through microscopy or flow cytometry. For microscopic analysis, cells were grown on custom glass-bottom cell culture plates and imaged under a Nikon Eclipse TE2000-U microscope controlled by MetaMorph software with a 100 \times objective lens. For flow cytometry studies, cells were detached with trypsin, washed, and analyzed.

DNA damage assessment

Oxidative DNA lesions were detected with an 8-oxoguanine antibody (Santa Cruz). Detached cells were fixed in 4% formaldehyde for 10 min at 37°C and permeabilized in ice-cold 90% methanol for 30 min on ice. After washing with PBS, cells were incubated in PBS containing anti-8-oxoguanine antibody (1:50), 0.5% Tween-20, and 5% FBS for 1 h. Cells were washed and incubated in PBS containing anti-mouse Texas Red antibody (1:500), 0.5% Tween-20, and 5% FBS for 1 h. Cells were then washed and analyzed with flow cytometer.

Mitochondrial DNA and mRNA measurement

Total cellular DNA was obtained using the DNeasy Blood & Tissue Kit (Qiagen) and used for detection with PCR primers listed below to quantify the nuclear (*18S rRNA*) to mitochondrial DNA (*CoI*) ratio as described (Brown and Clayton, 2002; Bai et al., 2004). Total cellular RNA was obtained using TRIzol (Life Technologies) and reverse transcribed to cDNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). PCR detection of expression genes was performed using the primers listed below comparing mitochondrial encoded oxidative phosphorylation genes (*ATP5 γ* and *CytC*) to nuclear encoded β -Actin as described (Yoon et al., 2010) (Table 2).

Western blot analysis

Cells were scraped from culture plates in the presence of RIPA buffer supplemented with Complete Protease Inhibitor (Roche,

TABLE 2. Primer sequences

Descriptions	Sequences
<i>CoI</i> F	5'-GCCCCAGATATAGCATTCCC-3'
<i>CoI</i> R	5'-GTTTCATCTGTTCTCTGCTCC-3'
<i>18S rRNA</i> F	5'-TAGAGGGACAAGTGGCGTTC-3'
<i>18S rRNA</i> R	5'-CGCTGAGCCAGTCAAGTGT-3'
<i>ATP5γ</i> F	5'-AGTTGGTGTGGCTGGATCA-3'
<i>ATP5γ</i> R	5'-GCTGCTTGAGAGATGGGTTC-3'
<i>CytC</i> F	5'-GGAGGCAAGCATAAGACTGG-3'
<i>CytC</i> R	5'-TCCATCAGGGTATCCTCTCC-3'
β -actin F	5'-TGTTACCAACTGGGACGACA-3'
β -actin R	5'-GGGGTGTGAAGGTCTCAA-3'

New York, NY), incubated on ice with frequent vortexing, and centrifuged. Supernatants were subjected to protein quantification and PAGE on 6–10% SDS gels followed by wet transfer to PVDF membranes and detection using β -catenin 1:1,000, CaV1.2 1:200, NF- κ B p65 1:200, and GAPDH 1:1,000 (Cell Signaling Technology, Danvers, MA).

Zebrafish

Adult wild-type AB zebrafish were obtained from the Zebrafish International Resource Center (Eugene, OR) and used for breeding. Embryos were injected with 1 mM antisense translation blocking morpholino oligos (MO; GeneTools) at the 1–2 cell stage. Zebrafish embryos were then cultured at 28.5°C in sterile egg water (Muntean et al., 2010). The following MO sequences were used: *control* MO: 5'-CCT CTT ACC TCA GTT ACA ATT TAT A-3', *cav1.2* MO: 5'-ACA TGT TTT TGC TTT CAT TTA CCA T-3', *pkd2* MO: 5'-AGG ACG AAC GCG ACT GGA GCT CAT C-3'. Knockdown of CaV1.2 was verified through Western blot analysis. Briefly, zebrafish embryos were dechorionated at 28 h postfertilization and homogenized in RIPA buffer to obtain protein extracts. Western was performed on 50 μ g total protein using CaV1.2 (1:200) and GAPDH (1:1,000) antibodies.

Histological examination was used to measure renal cyst formation and hydrocephalus at 3 days postfertilization. Embryos were fixed in a PBS solution containing 4% paraformaldehyde and 2% sucrose overnight at 4°C, dehydrated through an ethanol gradient, and embedded in JB4 resin (Polysciences, Inc., Warrington, PA) as specified in manufacturer's protocol. A Reichert Jung microtome was used to cut 5 μ m sections which were subsequently hematoxylin and eosin stained. Heart looping was assessed at 48 h postfertilization by positioning zebrafish on their dorsal axis and recording heart beat to reveal the respective relative locations of the atrium and the ventricle.

Data analysis

Data are reported as the mean \pm standard error of the mean. All image analysis was performed using ImageJ. All flow cytometry data were analyzed with BD Accuri C6 software and were presented without any compensation gating. All data were analyzed using IBM SPSS Statistics Version 21 software by performing the student t-test for two group comparison or ANOVA test followed by Tukey's post-test for three or more group comparison. Statistical significance is reported with a statistical power greater than 0.8 at $P < 0.05$.

Results

CaV1.2 is not required for primary cilia assembly

We recently reported that the voltage gated L-type calcium channel CaV1.2 localized to primary cilia in bovine LLC PK cells (Jin et al., 2013). We performed immunostaining to verify this finding in mouse renal epithelial cells (Fig. 1). The mouse *Tg737^{orp/orpk}* cell line contains a hypomorphic mutation in an intraflagellar transport gene (*Ift88*) that is required for cells




Fig. 1. Localization of CaV1.2 to renal epithelial cilia is not required for primary cilia assembly. Immunofluorescence revealed that CaV1.2 localized to primary cilia in renal epithelial cells (scrambled shRNA) when compared with cilia-deficient cells (*Tg737orp/orpk*). The presence of primary cilia was confirmed in CaV1.2 shRNA cells. Acetylated- α -tubulin was used as a ciliary marker. Arrow indicates the presence of primary cilium, except in cilia-deficient cells. Bar = 20 μ m.

to assemble primary cilia (Moyer et al., 1994). Thus, the *Tg737orp/orpk* system is a well-established model for studying cells without longer primary cilia, as verified through our immunostaining. We next asked if CaV1.2 played a role in primary cilia assembly. We generated a stable CaV1.2 shRNA knockdown mouse renal epithelial cell line and immunostaining studies revealed that primary cilia were similar to that of scrambled shRNA.

Wnt3a induces mitochondrial biogenesis in CaV1.2-deficient but not cilia-deficient cells

Wnt signaling has recently been reported to regulate mitochondrial physiology (Yoon et al., 2010). To assess mitochondrial mass, cells were stained with Mito Tracker Green (MTG) and observed live using fluorescence microscopy. When treated with recombinant Wnt3a, mitochondrial mass increased (Fig. 2a). However, the mitochondrial mass in *Tg737orp/orpk* cells was unchanged after Wnt3a treatment. We next performed this experiment in CaV1.2 shRNA cells and the results were similar to that of the scrambled control. To

quantify these findings, MTG fluorescence was recorded using flow cytometry which confirmed our fluorescent observation (Fig. 2b). Our MTG studies were further validated using a common technique by comparing mitochondrial DNA (*Coi*) to nuclear DNA (*18S rRNA*) (Brown and Clayton, 2002; Bai et al., 2004). As expected, Wnt3a did indeed statistically increase mitochondrial biogenesis in scrambled and CaV1.2 shRNA cells but not in *Tg737orp/orpk* cells (Fig. 2c). Our immunofluorescence study showed that Wnt3a did not alter CaV1.2 localization to cilia (Table 3).

Wnt3a increases mitochondrial activity in CaV1.2-deficient cells while decreasing activity in cilia-deficient cells

We next asked if Wnt3a would have an effect on mitochondrial oxidative phosphorylation (activity) in *Tg737orp/orpk* cells. Similar to before, we stained cells with Mito Tracker Red (MTR). Unlike MTG, MTR staining is dependent on the mitochondrial membrane potential (Poot and Pierce, 2001; Pendergrass et al., 2004). Therefore, increased staining




Fig. 2. Wnt3a induces mitochondrial biogenesis in *CaVI.2* shRNA but not *Tg737^{orp/orp}* cells. **a:** Mitochondrial mass was assessed by staining cells with Mito Tracker Green. Wnt3a was found to induce mitochondrial mass in scrambled and *CaVI.2* shRNA cells but had no effect on *Tg737^{orp/orp}* when examined using fluorescence microscopy (bar = 20 μ m). **b:** Results were quantified using flow cytometry. **c:** Mitochondrial DNA was measured using PCR by taking the ratio of a mitochondrial gene (*Coi*) to a nuclear gene (18S rRNA) (N = 3).

correlates to increased oxidative phosphorylation. As expected, Wnt3a increased MTR staining in scrambled and *CaVI.2* shRNA cells when observed using fluorescence microscopy (Fig. 3a). However, mitochondrial activity decreased in *Tg737^{orp/orp}* cells. We again quantified our findings using flow cytometry (Fig. 3b). Wnt3a significantly increased mitochondrial activity in scrambled and *CaVI.2* shRNA while significantly decreasing activity in *Tg737^{orp/orp}* cells. To verify these results, we compared expression of two key mitochondrial encoded oxidative phosphorylation genes (ATP Synthase 5 γ I and Cytochrome c) relative to that of nuclear encoded β -actin (Fig. 3c).

TABLE 3. *CaVI.2* ciliary localization

	% <i>CaVI.2</i> localization to cilia	N
PBS (control)		
Scramble shRNA	91.1	45
<i>Tg737^{orp/orp}</i>	91.7	36
<i>CaVI.2</i> shRNA	0.0	41
Wnt3a (100 ng/ml)		
Scramble shRNA	90.4	52
<i>Tg737^{orp/orp}</i>	92.3	39
<i>CaVI.2</i> shRNA	0.0	46

Wnt3a increases ROS and DNA damage in *CaVI.2*-deficient but not in cilia-deficient cells

An inevitable consequence of oxidative phosphorylation is the generation of reactive oxygen species (ROS) (Boveris et al., 1972; Boveris and Chance, 1973). MitoSOX is a cell permeable red fluorescent indicator specific for mitochondrial ROS. We therefore stained cells with MitoSOX and observed a significant increase in mitochondrial ROS in scrambled and *CaVI.2* shRNA after treatment with Wnt3a (Fig. 4a). A significant decrease in staining was observed in *Tg737^{orp/orp}* cells (Fig. 4b). Genomic DNA can be damaged by ROS to form DNA lesions resulting from mismatched repairs (Kasai et al., 1984). Thus, we quantified the levels of 8-Oxoguanine, a common DNA lesion formed by mismatched Adenine (Kasai, 1997). Treatment with Wnt3a was found to increase 8-Oxoguanine in scrambled and *CaVI.2* shRNA while no change was observed in *Tg737^{orp/orp}* cells (Fig. 4c).

Cilia modulates Wnt signaling to regulate *CaVI.2* expression

As previously reported, Wnt3a treatment induced β -catenin expression in all cells (Aberle et al., 1997). We confirmed this in our system, including in *Tg737^{orp/orp}* and *CaVI.2* shRNA cells (Fig. 5). Consistent with previous study (Corbit et al., 2008),




Fig. 3. Wnt3a increases mitochondrial activity in *CaV1.2* shRNA but decreasing activity in *Tg737orp/ork* cells. **a:** Mitochondrial oxidative phosphorylation was used to indicate activity through Mito Tracker Red staining. Wnt3a was found to increase oxidative phosphorylation in scrambled and *CaV1.2* shRNA cells while decreasing oxidative phosphorylation in *Tg737orp/ork* when examined using fluorescence microscopy (bar = 20 μ m). **b:** Results were quantified using flow cytometry. **c:** Two mitochondrial mRNAs encoded oxidative phosphorylation genes (CytC and ATP5 γ 1) were measured using PCR normalized to nuclear encoded β -actin (N = 3).

Tg737orp/ork cells showed a higher basal level of β -catenin than control. Further, Wnt3a treatment increased *CaV1.2* expression in scrambled shRNA while decreasing *CaV1.2* in *Tg737orp/ork* cells. Of note is that *CaV1.2* expression was not detectable in *CaV1.2* shRNA cells, confirming knockdown of *CaV1.2* in our stable cell line.

The DNA damage response (DDR) is a cellular mechanism to recover from DNA lesions, such as 8-Oxoguanine (Kasai et al., 1984; Jackson and Bartek, 2009). One arm of this cell survival pathway is the activation of nuclear factor κ B p65 (NF κ B p65) (Janssens and Tschopp, 2006). Through Western blot analysis, we also found that Wnt3a induced NF κ B p65 expression in scrambled and *CaV1.2* shRNA (Fig. 5). On the other hand, *Tg737orp/ork* cells expressed a high basal level of NF κ B p65 which decreased in response to Wnt3a. In addition, *CaV1.2* expression was found to correlate with NF κ B p65.

CaV1.2 knockdown zebrafish develop PKD phenotypes

We have shown that *CaV1.2* localizes to primary cilia and have now elucidated the mechanism by which *CaV1.2* expression is regulated in renal epithelial cells. To assess the biological significance of *CaV1.2* expression, we used antisense

morpholinos to knockdown *CaV1.2* in zebrafish. Knockdown of the ciliary calcium channel polycystin-2 in zebrafish has been reported to result in PKD phenotypes including renal cyst formation, hydrocephalus, and left-right asymmetry (Obara et al., 2006). Our study showed that knockdown of *pkd2* increased *CaV1.2* expression (Fig. 6). This slight increase in *CaV1.2* was significant compared to the control morpholino. Interestingly, similar phenotypes were observed in *CaV1.2* morpholino (*cav1.2* MO) zebrafish. Compared with a non-specific control morpholino (control MO) injection, *cav1.2* MO zebrafish developed renal cysts (Fig. 7a), hydrocephalus (Fig. 7b) and various heart-looping defects (Fig. 7c). As generally accepted (Bakkers, 2011), left-right asymmetry was assessed by measuring the relative position of the cardiac atrium and ventricle with respect to the dorsal axis (Supplemental Movie 1).

Discussion

Non-motile primary cilia have been found to play a critical role in Wnt signaling by restricting β -catenin accumulation. Overexpression of polycystin-1 (a ciliary signaling receptor) inhibits GSK3 β and stabilizes β -catenin (Kim et al., 1999).




Fig. 4. Wnt3a increases ROS and DNA damage in *CaV1.2* shRNA but not in *Tg737orp/orpk* cells. **a:** Mitochondrial ROS was assessed by staining cells with MitoSOX. Wnt3a was found to increase ROS in scrambled and *CaV1.2* shRNA cells while decreasing ROS in *Tg737orp/orpk* cells when examined using fluorescence microscopy (bar = 20 μ m). **b:** Results were quantified using flow cytometry. **c:** Wnt3a increased formation of the oxidative DNA lesion 8-Oxoguanine in scrambled and *CaV1.2* shRNA cells but had no effect on *Tg737orp/orpk* cells (N = 3).

Polycystin-2 (encoded by *Pkd2*) is calcium channel forming protein found in primary cilia. In *Pkd2*^{-/-} embryos, cilia length was found to be decreased while β -catenin was upregulated (Kim et al., 2009). Interestingly, transgenic mice overexpressing β -catenin also developed cystic kidneys (Saadi-Kheddouci et al., 2001). Further, *LRP6*^{-/-} (a component of the Wnt receptor complex) mouse embryos die in utero with cystic kidneys (Pinson et al., 2000). Thus, primary cilia and Wnt signaling play a crucial role in PKD (Corbit et al., 2008).

Given that Wnt signaling also modulates mitochondrial physiology (Yoon et al., 2010), we examined the role of cilia in regards to mitochondria. *Tg737orp/orpk* contains an intron insertion at the 3' end of the intraflagellar transport 88 (*Ift88*) gene which results in a hypomorphic mutation that prevents ciliogenesis (Moyer et al., 1994). We used *Tg737orp/orpk* cells as a model for a cilia-deficient system. Through immunostaining, we confirmed the absence of cilia in *Tg737orp/orpk* cells compared with control (Fig. 1). The voltage-gated L-type calcium channel *CaV1.2* also localized to primary cilia. We generated a stable *CaV1.2* shRNA cell line and observed no changes in primary cilia compared with control. Further, treatment with Wnt3a had no effect on cilia number or length in scrambled or *CaV1.2* shRNA cells (data not shown). Thus, *CaV1.2* does not seem to play a role in ciliogenesis.

Mitochondrial biogenesis, oxidative phosphorylation, and generation of reactive oxidative species (ROS) were increased in response to Wnt3a in control renal epithelial cells

(Figures 2–4). The elevated levels of oxidative stress increased the formation of DNA lesions and the cellular DNA damage response (DDR). An interesting aspect of this response was an increase in expression of *CaV1.2* (Fig. 5). In *CaV1.2* knockdown cells, Wnt3a induced a similar effect on mitochondria and DDR. This data suggests that *CaV1.2* is a downstream effector in regard to Wnt signaling. In cilia-deficient cells, Wnt3a was unable to induce mitochondrial biogenesis and decreased mitochondrial activity, ROS production, and DDR. *CaV1.2* was found to be overexpressed in cilia-deficient cells as a compensatory mechanism; however, its expression decreased following Wnt3a treatment. Therefore, cilia length plays a role in regulating *CaV1.2* expression through modulation of Wnt signaling.

Defective primary cilia, indicated by either depletion of key ciliary proteins or fundamental changes in structure/length, results in PKD phenotypes (Wilson, 2004). Here we show that *CaV1.2* is a biologically significant ciliary protein. In the absence of *CaV1.2* in zebrafish (Fig. 6), PKD phenotypes including renal cyst formation, hydrocephalus, and left-right asymmetry defects were observed (Fig. 7). Moreover, *CaV1.2* was found to be overexpressed in *pkd2* knockdown zebrafish. This is intriguing given that both *CaV1.2* and PC2 are calcium channel forming proteins in the primary cilium, which further suggests a role for *CaV1.2* in PKD pathogenesis. Therefore, *CaV1.2* not only localizes to renal epithelial primary cilia, but it is also required for cilia function.




Fig. 5. Cilia modulates Wnt signaling to regulate CaV1.2 expression. Western blotting was performed on cellular protein extracts. Wnt3a treatment increased β-catenin accumulation in all cells, however, *Tg737^{orp/orp}* cells displayed high basal levels of β-catenin. NF-κB p65 was blotted as a readout for DNA damage response (DDR) to ROS induced DNA lesions. Wnt3a induced NF-κB p65 expression in scrambled and *CaV1.2* shRNA cells but decreased NF-κB p65 in *Tg737^{orp/orp}* cells. *CaV1.2* was overexpressed in *Tg737^{orp/orp}* cells. Wnt3a treatment induced *CaV1.2* expression in scrambled shRNA while decreasing expression in *Tg737^{orp/orp}* cells. Data normalized to GAPDH for analysis (N = 3). Asterisks indicate significant difference from the corresponding control group ($P < 0.05$). # signs denote significant difference from the corresponding scrambled shRNA group ($P < 0.05$).




Fig. 6. *pkd2* knockdown increases CaV1.2 expression in zebrafish. Embryonic zebrafish protein extracts were obtained at 28 h postfertilization. Western blotting showed that *cav1.2* MO effectively reduced CaV1.2 expression compared with control MO. *pkd2* MO zebrafish were used to further verify morpholino specificity. Results were quantified through one-way ANOVA with Tukey post-test. Statistical significance is reported with a mean difference at the 0.05 level and denoted with an asterisk (N = 3).




a renal cyst**b hydrocephalus****c heart looping**

Fig. 7. CaV1.2 knockdown zebrafish develop renal cysts, hydrocephalus, and left-right asymmetry defects. CaV1.2 was knocked down in zebrafish (cav1.2 MO) using morpholino microinjection and compared to a scrambled control morpholino (control MO) to assess phenotypes. Renal cyst formation (a) and hydrocephalus (b) were measured at 3 days postfertilization through standard H&E staining as indicated by arrows. Left-right asymmetry was determined by measuring heart looping (c) at 2 days postfertilization under live microscopy.

Although it has been known that abnormal Wnt signaling leads to renal cyst formation (Lancaster et al., 2009), the underlying mechanism has been unclear. One proposed explanation is that Wnt signaling regulates renal cell proliferation and planar cell polarity to maintain renal tubule homeostasis (Happe et al., 2009). Our present study further suggests that cilia regulate Wnt signaling which ultimately controls CaV1.2 expression. Therefore, in the absence of CaV1.2, ciliary function is compromised leading to formation of renal cysts. In addition, our study also shows a previously unrecognized relationship between primary cilia and mitochondrial function. Overall, we propose that Wnt3a induces β -catenin accumulation in a cilia-dependent manner. This in turn increases mitochondrial biogenesis, oxidative phosphorylation, and generation of ROS that cause genomic DNA damage. The DDR triggers NF κ B p65 expression ultimately resulting in increased CaV1.2 expression.

Acknowledgments

This work was supported in part by DOD PR130153 and NIH DK080640.

Literature Cited

- Aberle H, Bauer A, Stappert J, Kispert A, Kemler R. 1997. Beta-catenin is a target for the ubiquitin-proteasome pathway. *EMBO J* 16:3797–3804.
- AbouAlaiwi WA, Takahashi M, Mell BR, Jones TJ, Ratnam S, Kolb RJ, Nauli SM. 2009. Ciliary polycystin-2 is a mechanosensitive calcium channel involved in nitric oxide signaling cascades. *Circ Res* 104:860–869.

- AbouAlaiwi WA, Muntean BS, Ratnam S, Joe B, Liu L, Booth RL, Rodriguez I, Herbert BS, Bacallao RL, Fruttiger M, Mak TW, Zhou J, Nauli SM. 2013. Survivin-induced abnormal ploidy contributes to cystic kidney and aneurysm formation. *Circulation*.
- Bai RK, Pong CL, Hsu CH, Wong LJ. 2004. Quantitative PCR analysis of mitochondrial DNA content in patients with mitochondrial disease. *Ann NY Acad Sci* 1011:304–309.
- Bakkers J. 2011. Zebrafish as a model to study cardiac development and human cardiac disease. *Cardiovasc Res* 91:279–288.
- Boveris A, Chance B. 1973. The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem J* 134:707–716.
- Boveris A, Oshino N, Chance B. 1972. The cellular production of hydrogen peroxide. *Biochem J* 128:617–630.
- Brown TA, Clayton DA. 2002. Release of replication termination controls mitochondrial DNA copy number after depletion with 2',3'-dideoxycytidine. *Nucleic acids Res* 30:2004–2010.
- Carter CS, Vogel TW, Zhang Q, Seo S, Swiderski RE, Moninger TO, Cassell MD, Thedens DR, Keppler-Noreuil KM, Nopoulos P, Nishimura DY, Searby CC, Bugge K, Sheffield VC. 2012. Abnormal development of NG2+PDGFR- α + neural progenitor cells leads to neonatal hydrocephalus in a ciliopathy mouse model. *Nat Med* 18:1797–1804.
- Corbit KC, Shyer AE, Dowdle WE, Gauden J, Singla V, Chen MH, Chuang PT, Reiter JF. 2008. Kif3a constrains beta-catenin-dependent Wnt signalling through dual ciliary and non-ciliary mechanisms. *Nat Cell Biol* 10:70–76.
- Gerdes JM, Liu Y, Zaghloul NA, Leitch CC, Lawson SS, Kato M, Beachy PA, Beales PL, DeMartino GN, Fisher S, Badano JL, Katsanis N. 2007. Disruption of the basal body compromises proteasomal function and perturbs intracellular Wnt response. *Nat Genet* 39:1350–1360.
- Happe H, Leonhard WN, van der Wal A, van de Water B, Lantinga-van Leeuwen IS, Breuning MH, de Heer E, Peters DJ. 2009. Toxic tubular injury in kidneys from Pkd1-deletion mice accelerates cystogenesis accompanied by dysregulated planar cell polarity and canonical Wnt signaling pathways. *Hum Mol Genet* 18:2532–2542.
- Jackson SP, Bartek J. 2009. The DNA-damage response in human biology and disease. *Nature* 461:1071–1078.
- Janssens S, Tschopp J. 2006. Signals from within: The DNA-damage-induced NF- κ B response. *Cell Death Differ* 13:773–784.
- Jin X, Mohieldin AM, Muntean BS, Green JA, Shah JV, Mykityn K, Nauli SM. 2013. Cilioplasm is a cellular compartment for calcium signaling in response to mechanical and chemical stimuli. *Cell Mol Life Sci*.
- Kasai H. 1997. Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis. *Mutat Res* 387:147–163.
- Kasai H, Tanooka H, Nishimura S. 1984. Formation of 8-hydroxyguanine residues in DNA by X-irradiation. *Gann* 75:1037–1039.

- Kim E, Arnould T, Sellin LK, Benzing T, Fan MJ, Gruning W, Sokol SY, Drummond I, Walz G. 1999. The polycystic kidney disease 1 gene product modulates Wnt signaling. *J Biol Chem* 274:4947–4953.
- Kim I, Ding T, Fu Y, Li C, Cui L, Li A, Lian P, Liang D, Wang DW, Guo C, Ma J, Zhao P, Coffey RJ, Zhan Q, Wu G. 2009. Conditional mutation of Pkd2 causes cystogenesis and upregulates beta-catenin. *J Am Soc Nephrol* 20:2556–2569.
- Lancaster MA, Louie CM, Silhavy JL, Sintasath L, Decambre M, Nigam SK, Willert K, Gleeson JG. 2009. Impaired Wnt-beta-catenin signaling disrupts adult renal homeostasis and leads to cystic kidney ciliopathy. *Nat Med* 15:1046–1054.
- Lancaster MA, Schroth J, Gleeson JG. 2011. Subcellular spatial regulation of canonical Wnt signalling at the primary cilium. *Nat Cell Biol* 13:700–707.
- Moyer JH, Lee-Tischler MJ, Kwon HY, Schrick JJ, Avner ED, Sweeney WE, Godfrey VL, Cacheiro NL, Wilkinson JE, Woychik RP. 1994. Candidate gene associated with a mutation causing recessive polycystic kidney disease in mice. *Science* 264:1329–1333.
- Muntean BS, Horvat CM, Behler JH, Aboulaiwi WA, Nauli AM, Williams FE, Nauli SM. 2010. A comparative study of embedded and anesthetized zebrafish in vivo on myocardial calcium oscillation and heart muscle contraction. *Front Pharmacol* 1:139.
- Muntean BS, Jin X, Nauli SM. Chapter 9: Primary cilia are mechanosensory organelles with chemosensory roles. *Mechanosensitivity and mechanotransduction*. ISBN: 978-994-007-2003-2009.
- Nauli SM, Alenghat FJ, Luo Y, Williams E, Vassilev P, Li X, Elia AE, Lu W, Brown EM, Quinn SJ, Ingber DE, Zhou J. 2003. Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet* 33:129–137.
- Nauli SM, Kawanabe Y, Kaminski JJ, Pearce WJ, Ingber DE, Zhou J. 2008. Endothelial cilia are fluid shear sensors that regulate calcium signaling and nitric oxide production through polycystin-1. *Circulation* 117:1161–1171.
- Norris DP. 2012. Cilia, calcium and the basis of left-right asymmetry. *BMC Biol* 10:102.
- Obara T, Mangos S, Liu Y, Zhao J, Wiessner S, Kramer-Zucker AG, Olale F, Schier AF, Drummond IA. 2006. Polycystin-2 immunolocalization and function in zebrafish. *J Am Soc Nephrol* 17:2706–2718.
- Pendergrass W, Wolf N, Poot M. 2004. Efficacy of MitoTracker Green and CMXRosamine to measure changes in mitochondrial membrane potentials in living cells and tissues. *Cytometry A* 61:162–169.
- Pinson KI, Brennan J, Monkley S, Avery BJ, Skarnes WC. 2000. An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature* 407:535–538.
- Poot M, Pierce RH. 2001. Analysis of mitochondria by flow cytometry. *Methods Cell Biol* 64:117–128.
- Saadi-Kheddouci S, Berrebi D, Romagnolo B, Cluzeaud F, Peuchmaur M, Kahn A, Vandewalle A, Perret C. 2001. Early development of polycystic kidney disease in transgenic mice expressing an activated mutant of the beta-catenin gene. *Oncogene* 20:5972–5981.
- Wilson PD. 2004. Polycystic kidney disease. *N Engl J Med* 350:151–164.
- Yoon JC, Ng A, Kim BH, Bianco A, Xavier RJ, Elledge SJ. 2010. Wnt signaling regulates mitochondrial physiology and insulin sensitivity. *Genes Dev* 24:1507–1518.

Supporting Information

Additional supporting information may be found in the online version of this article at the publisher's web-site.

Primary Cilium Regulates CaV1.2 Expression Through Wnt Signaling

BRIAN S. MUNTEAN,¹ XINGJIAN JIN,² FREDERICK E. WILLIAMS,² AND SURYA M. NAULI^{1,2*}

¹Department of Medicinal and Biological Chemistry, The University of Toledo, Toledo, Ohio

²Department of Pharmacology, The University of Toledo, Toledo, Ohio

Primary cilia are sensory organelles that provide a feedback mechanism to restrict Wnt signaling in the absence of endogenous Wnt activators. Abnormal Wnt signaling has been shown to result in polycystic kidney disease (PKD) although the exact mechanism has been debated. Previously, we reported that the calcium channel CaV1.2 functions in primary cilia. In this study, we show that CaV1.2 expression level is regulated by Wnt signaling. This occurs through modulation of mitochondrial mass and activity resulting in increased reactive oxygen species which generate oxidative DNA lesions. We found that the subsequent cellular DNA damage response triggers increased CaV1.2 expression. In the absence of primary cilia where Wnt signaling is upregulated, we found that CaV1.2 is overexpressed as a compensatory mechanism. We show for the first time that CaV1.2 knockdown in zebrafish results in classic primary cilia defects including renal cyst formation, hydrocephalus, and left-right asymmetry defects. Our study shows that suppressed Wnt signaling prevents CaV1.2 expression ultimately resulting in PKD phenotypes. Thus, CaV1.2 expression is tightly regulated through Wnt signaling and plays an essential sensory role in primary cilia necessary for cellular homeostasis.

J. Cell. Physiol. 229: 1926–1934, 2014. © 2014 Wiley Periodicals, Inc.

Wnt signaling is an important regulator of cellular development and proliferation. In the absence of Wnt ligands, a complex consisting of Axin, adenomatous polyposis coli (APC), and glycogen synthase kinase 3 β (GSK3 β) induces β -catenin for ubiquitylation by SCF E3 ligases and eventual proteasomal degradation (Aberle et al., 1997). Wnt signal transduction occurs when secreted Wnt ligands bind Frizzled receptors resulting in phosphorylation of LRP5/6. The Axin-APC-GSK3 β complex is then recruited to LRP5/6 at the cell membrane which prevents β -catenin from being degraded. The accumulated β -catenin translocates to the nucleus and activates transcription of Wnt target genes (Muntean et al., 2012).

Primary cilia are non-motile sensory organelles present as a single copy on most differentiated cells in the body. Calcium signaling through primary cilium is essential for renal epithelial homeostasis (Nauli et al., 2003; Jin et al., 2013). Cilia extend from the cell surface through the basal body via intraflagellar transport (Moyer et al., 1994). The most common pathologies resulting from cilia dysfunction include polycystic kidney (Wilson, 2004), hypertension (Nauli et al., 2008; AbouAlaiwi et al., 2009), aneurysm (AbouAlaiwi et al., 2013), hydrocephalus (Carter et al., 2012), and left-right asymmetry defects (Norris, 2012).

Abnormal Wnt signaling has also been linked to polycystic kidney disease (PKD) (Lancaster et al., 2009). For example, increased cytosolic and nuclear β -catenin accumulation has been shown in various cilia mutant cells (Gerdes et al., 2007; Lancaster et al., 2011). Thus, primary cilia are thought to provide a feedback mechanism that restricts Wnt signaling in the absence of appropriate ligands (Gerdes et al., 2007; Lancaster et al., 2009, 2011).

We recently showed that voltage-gated L-type calcium channel CaV1.2 localized to primary cilia in renal epithelia (Jin et al., 2013). Because Wnt signaling has also been reported to modulate mitochondrial physiology (Yoon et al., 2010), we hypothesized that primary cilia play a role in Wnt regulation of mitochondria through CaV1.2. We show that although CaV1.2 is not required for cilia formation, Wnt increases mitochondria mass and activity in CaV1.2 deficient renal epithelial cells. This increases mitochondria reactive oxidative species (ROS) and DNA damage, resulting in PKD phenotypes. Thus, our study suggests that primary cilia may play a role in CaV1.2 expression level through Wnt regulation of mitochondria.

Materials and Methods

The experimental use of zebrafish was approved by The University of Toledo's Institutional Animal Care and Use Committee (IACUC). The use of lentiviral components was approved by the Institutional Biosafety Committee of The University of Toledo.

Cell culture

Immortalized mouse renal epithelial wild-type and *Tg737^{orp/k}* cells were cultured in Dulbecco's Modified Eagle Medium (Corning Cellgro) supplemented with 10% fetal bovine serum (HyClone Laboratories, Logan, Utah) and 1% penicillin/streptomycin (Corning Cellgro) at 39°C in 5% CO₂, as previously described (AbouAlaiwi et al., 2013). Prior to experiments, cells were treated with 100 ng/ml recombinant Wnt3a (R&D Systems, Minneapolis, MN) for 3 days and serum starved for 24 h.

RNAi knockdown cells

shRNA lentiviral vectors specific to *Cacna1c* (Origene; pGFP-C-shLenti clone ID: TL500242) were transfected into HEK293T cells. Viral supernatants were collected after 48 h, centrifuged, and passed through a 0.45 μ m filter. Cells were then spin-inoculated with pseudoviral particles containing 8 μ g/ml polybrene at 2,500 rpm for 30 min at 30°C and then cultured for up to 1 week. CaV1.2 knockdown was verified through Western blot analysis.

Brian S. Muntean and Xingjian Jin contributed equally to this work.

* Correspondence to: Surya M. Nauli, Department of Pharmacology, The University of Toledo, MS 1015, Health Education Building; Room 274, 3000 Arlington Ave, Toledo, OH 43614.

E-mail: surya.nauli@utoledo.edu

Manuscript Received: 20 January 2014

Manuscript Accepted: 1 April 2014

Accepted manuscript online in Wiley Online Library (wileyonlinelibrary.com): 2 April 2014.

DOI: 10.1002/jcp.24642

TABLE 1. shRNA sequences

Descriptions	Sequences
Scrambled control	5'-TGACCACCCTGACCTACGGCGTGCAGTGC-3'
<i>Cacna1c</i>	5'-TCAGAGTGCCTCACTGTTCTCGTGACCT-3'

Stable knockdown cell lines were obtained through puromycin selection. The following shRNA sequences were used (Table 1).

Immunostaining studies

Cells were grown to confluence on collagen-coated glass coverslips and differentiated in serum-free media for 24 h. Cells were then fixed in 4% paraformaldehyde in PBS containing 2% sucrose, permeabilized in 10% triton X-100, incubated sequentially with primary followed by secondary antibodies for 1 h each, and finally mounted on a slide with DAPI hard set mounting media (Vector Laboratories, Burlingame, CA). The following primary antibody dilutions were used: acetylated- α -tubulin 1:10,000 (Sigma-Aldrich, St. Louis, MO) and CaV1.2 1:50 (Alomone Labs, Jerusalem, Israel). Anti-mouse Texas Red and anti-rabbit FITC fluorescent conjugated secondary antibodies were used at 1:500 (VectorLabs).

Mitochondrial studies

MitoTracker Green FM and MitoTracker Red CMXRos (Cell Signaling Technology) were incubated with cells at 100 nM for 30 min at 37°C. MitoSOX (Life Technologies) was incubated with cells at 5 μ M for 10 min at 37°C. After staining, cells were washed three times with PBS and analyzed immediately through microscopy or flow cytometry. For microscopic analysis, cells were grown on custom glass-bottom cell culture plates and imaged under a Nikon Eclipse TE2000-U microscope controlled by MetaMorph software with a 100 \times objective lens. For flow cytometry studies, cells were detached with trypsin, washed, and analyzed.

DNA damage assessment

Oxidative DNA lesions were detected with an 8-oxoguanine antibody (Santa Cruz). Detached cells were fixed in 4% formaldehyde for 10 min at 37°C and permeabilized in ice-cold 90% methanol for 30 min on ice. After washing with PBS, cells were incubated in PBS containing anti-8-oxoguanine antibody (1:50), 0.5% Tween-20, and 5% FBS for 1 h. Cells were washed and incubated in PBS containing anti-mouse Texas Red antibody (1:500), 0.5% Tween-20, and 5% FBS for 1 h. Cells were then washed and analyzed with flow cytometer.

Mitochondrial DNA and mRNA measurement

Total cellular DNA was obtained using the DNeasy Blood & Tissue Kit (Qiagen) and used for detection with PCR primers listed below to quantify the nuclear (*18S rRNA*) to mitochondrial DNA (*CoI*) ratio as described (Brown and Clayton, 2002; Bai et al., 2004). Total cellular RNA was obtained using TRIzol (Life Technologies) and reverse transcribed to cDNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). PCR detection of expression genes was performed using the primers listed below comparing mitochondrial encoded oxidative phosphorylation genes (*ATP5 γ* and *CytC*) to nuclear encoded β -Actin as described (Yoon et al., 2010) (Table 2).

Western blot analysis

Cells were scraped from culture plates in the presence of RIPA buffer supplemented with Complete Protease Inhibitor (Roche,

TABLE 2. Primer sequences

Descriptions	Sequences
<i>CoI</i> F	5'-GCCCCAGATATAGCATTCCC-3'
<i>CoI</i> R	5'-GTTTCATCTGTTCTCTGCTCC-3'
<i>18S rRNA</i> F	5'-TAGAGGGACAAGTGGCGTTC-3'
<i>18S rRNA</i> R	5'-CGCTGAGCCAGTCAAGTGT-3'
<i>ATP5γ</i> F	5'-AGTTGGTGTGGCTGGATCA-3'
<i>ATP5γ</i> R	5'-GCTGCTTGAGAGATGGGTTC-3'
<i>CytC</i> F	5'-GGAGGCAAGCATAAGACTGG-3'
<i>CytC</i> R	5'-TCCATCAGGGTATCCTCTCC-3'
β -actin F	5'-TGTACCAACTGGGACGACA-3'
β -actin R	5'-GGGGTGTGAAGGTCTCAA-3'

New York, NY), incubated on ice with frequent vortexing, and centrifuged. Supernatants were subjected to protein quantification and PAGE on 6–10% SDS gels followed by wet transfer to PVDF membranes and detection using β -catenin 1:1,000, CaV1.2 1:200, NF- κ B p65 1:200, and GAPDH 1:1,000 (Cell Signaling Technology, Danvers, MA).

Zebrafish

Adult wild-type AB zebrafish were obtained from the Zebrafish International Resource Center (Eugene, OR) and used for breeding. Embryos were injected with 1 mM antisense translation blocking morpholino oligos (MO; GeneTools) at the 1–2 cell stage. Zebrafish embryos were then cultured at 28.5°C in sterile egg water (Muntean et al., 2010). The following MO sequences were used: *control* MO: 5'-CCT CTT ACC TCA GTT ACA ATT TAT A-3', *cav1.2* MO: 5'-ACA TGT TTT TGC TTT CAT TTA CCA T-3', *pkd2* MO: 5'-AGG ACG AAC GCG ACT GGA GCT CAT C-3'. Knockdown of CaV1.2 was verified through Western blot analysis. Briefly, zebrafish embryos were dechorionated at 28 h postfertilization and homogenized in RIPA buffer to obtain protein extracts. Western was performed on 50 μ g total protein using CaV1.2 (1:200) and GAPDH (1:1,000) antibodies.

Histological examination was used to measure renal cyst formation and hydrocephalus at 3 days postfertilization. Embryos were fixed in a PBS solution containing 4% paraformaldehyde and 2% sucrose overnight at 4°C, dehydrated through an ethanol gradient, and embedded in JB4 resin (Polysciences, Inc., Warrington, PA) as specified in manufacturer's protocol. A Reichert Jung microtome was used to cut 5 μ m sections which were subsequently hematoxylin and eosin stained. Heart looping was assessed at 48 h postfertilization by positioning zebrafish on their dorsal axis and recording heart beat to reveal the respective relative locations of the atrium and the ventricle.

Data analysis

Data are reported as the mean \pm standard error of the mean. All image analysis was performed using ImageJ. All flow cytometry data were analyzed with BD Accuri C6 software and were presented without any compensation gating. All data were analyzed using IBM SPSS Statistics Version 21 software by performing the student t-test for two group comparison or ANOVA test followed by Tukey's post-test for three or more group comparison. Statistical significance is reported with a statistical power greater than 0.8 at $P < 0.05$.

Results

CaV1.2 is not required for primary cilia assembly

We recently reported that the voltage gated L-type calcium channel CaV1.2 localized to primary cilia in bovine LLC PK cells (Jin et al., 2013). We performed immunostaining to verify this finding in mouse renal epithelial cells (Fig. 1). The mouse *Tg737^{orp/orpk}* cell line contains a hypomorphic mutation in an intraflagellar transport gene (*Ift88*) that is required for cells




Fig. 1. Localization of CaV1.2 to renal epithelial cilia is not required for primary cilia assembly. Immunofluorescence revealed that CaV1.2 localized to primary cilia in renal epithelial cells (scrambled shRNA) when compared with cilia-deficient cells (*Tg737^{orp/orpk}*). The presence of primary cilia was confirmed in CaV1.2 shRNA cells. Acetylated- α -tubulin was used as a ciliary marker. Arrow indicates the presence of primary cilium, except in cilia-deficient cells. Bar = 20 μ m.

to assemble primary cilia (Moyer et al., 1994). Thus, the *Tg737^{orp/orpk}* system is a well-established model for studying cells without longer primary cilia, as verified through our immunostaining. We next asked if CaV1.2 played a role in primary cilia assembly. We generated a stable CaV1.2 shRNA knockdown mouse renal epithelial cell line and immunostaining studies revealed that primary cilia were similar to that of scrambled shRNA.

Wnt3a induces mitochondrial biogenesis in CaV1.2-deficient but not cilia-deficient cells

Wnt signaling has recently been reported to regulate mitochondrial physiology (Yoon et al., 2010). To assess mitochondrial mass, cells were stained with Mito Tracker Green (MTG) and observed live using fluorescence microscopy. When treated with recombinant Wnt3a, mitochondrial mass increased (Fig. 2a). However, the mitochondrial mass in *Tg737^{orp/orpk}* cells was unchanged after Wnt3a treatment. We next performed this experiment in CaV1.2 shRNA cells and the results were similar to that of the scrambled control. To

quantify these findings, MTG fluorescence was recorded using flow cytometry which confirmed our fluorescent observation (Fig. 2b). Our MTG studies were further validated using a common technique by comparing mitochondrial DNA (*Coi*) to nuclear DNA (*18S rRNA*) (Brown and Clayton, 2002; Bai et al., 2004). As expected, Wnt3a did indeed statistically increase mitochondrial biogenesis in scrambled and CaV1.2 shRNA cells but not in *Tg737^{orp/orpk}* cells (Fig. 2c). Our immunofluorescence study showed that Wnt3a did not alter CaV1.2 localization to cilia (Table 3).

Wnt3a increases mitochondrial activity in CaV1.2-deficient cells while decreasing activity in cilia-deficient cells

We next asked if Wnt3a would have an effect on mitochondrial oxidative phosphorylation (activity) in *Tg737^{orp/orpk}* cells. Similar to before, we stained cells with Mito Tracker Red (MTR). Unlike MTG, MTR staining is dependent on the mitochondrial membrane potential (Poot and Pierce, 2001; Pendergrass et al., 2004). Therefore, increased staining




Fig. 2. Wnt3a induces mitochondrial biogenesis in *CaVI.2* shRNA but not *Tg737^{orp/orp}* cells. **a:** Mitochondrial mass was assessed by staining cells with Mito Tracker Green. Wnt3a was found to induce mitochondrial mass in scrambled and *CaVI.2* shRNA cells but had no effect on *Tg737^{orp/orp}* when examined using fluorescence microscopy (bar = 20 μ m). **b:** Results were quantified using flow cytometry. **c:** Mitochondrial DNA was measured using PCR by taking the ratio of a mitochondrial gene (*Coi*) to a nuclear gene (18S rRNA) (N = 3).

correlates to increased oxidative phosphorylation. As expected, Wnt3a increased MTR staining in scrambled and *CaVI.2* shRNA cells when observed using fluorescence microscopy (Fig. 3a). However, mitochondrial activity decreased in *Tg737^{orp/orp}* cells. We again quantified our findings using flow cytometry (Fig. 3b). Wnt3a significantly increased mitochondrial activity in scrambled and *CaVI.2* shRNA while significantly decreasing activity in *Tg737^{orp/orp}* cells. To verify these results, we compared expression of two key mitochondrial encoded oxidative phosphorylation genes (ATP Synthase 5 γ I and Cytochrome c) relative to that of nuclear encoded β -actin (Fig. 3c).

TABLE 3. *CaVI.2* ciliary localization

	% <i>CaVI.2</i> localization to cilia	N
PBS (control)		
Scramble shRNA	91.1	45
<i>Tg737^{orp/orp}</i>	91.7	36
<i>CaVI.2</i> shRNA	0.0	41
Wnt3a (100 ng/ml)		
Scramble shRNA	90.4	52
<i>Tg737^{orp/orp}</i>	92.3	39
<i>CaVI.2</i> shRNA	0.0	46

Wnt3a increases ROS and DNA damage in *CaVI.2*-deficient but not in cilia-deficient cells

An inevitable consequence of oxidative phosphorylation is the generation of reactive oxygen species (ROS) (Boveris et al., 1972; Boveris and Chance, 1973). MitoSOX is a cell permeable red fluorescent indicator specific for mitochondrial ROS. We therefore stained cells with MitoSOX and observed a significant increase in mitochondrial ROS in scrambled and *CaVI.2* shRNA after treatment with Wnt3a (Fig. 4a). A significant decrease in staining was observed in *Tg737^{orp/orp}* cells (Fig. 4b). Genomic DNA can be damaged by ROS to form DNA lesions resulting from mismatched repairs (Kasai et al., 1984). Thus, we quantified the levels of 8-Oxoguanine, a common DNA lesion formed by mismatched Adenine (Kasai, 1997). Treatment with Wnt3a was found to increase 8-Oxoguanine in scrambled and *CaVI.2* shRNA while no change was observed in *Tg737^{orp/orp}* cells (Fig. 4c).

Cilia modulates Wnt signaling to regulate *CaVI.2* expression

As previously reported, Wnt3a treatment induced β -catenin expression in all cells (Aberle et al., 1997). We confirmed this in our system, including in *Tg737^{orp/orp}* and *CaVI.2* shRNA cells (Fig. 5). Consistent with previous study (Corbit et al., 2008),




Fig. 3. Wnt3a increases mitochondrial activity in *CaV1.2* shRNA but decreasing activity in *Tg737orp/ork* cells. **a:** Mitochondrial oxidative phosphorylation was used to indicate activity through Mito Tracker Red staining. Wnt3a was found to increase oxidative phosphorylation in scrambled and *CaV1.2* shRNA cells while decreasing oxidative phosphorylation in *Tg737orp/ork* when examined using fluorescence microscopy (bar = 20 μ m). **b:** Results were quantified using flow cytometry. **c:** Two mitochondrial mRNAs encoded oxidative phosphorylation genes (CytC and ATP5 γ 1) were measured using PCR normalized to nuclear encoded β -actin (N = 3).

Tg737orp/ork cells showed a higher basal level of β -catenin than control. Further, Wnt3a treatment increased *CaV1.2* expression in scrambled shRNA while decreasing *CaV1.2* in *Tg737orp/ork* cells. Of note is that *CaV1.2* expression was not detectable in *CaV1.2* shRNA cells, confirming knockdown of *CaV1.2* in our stable cell line.

The DNA damage response (DDR) is a cellular mechanism to recover from DNA lesions, such as 8-Oxoguanine (Kasai et al., 1984; Jackson and Bartek, 2009). One arm of this cell survival pathway is the activation of nuclear factor κ B p65 (NF κ B p65) (Janssens and Tschopp, 2006). Through Western blot analysis, we also found that Wnt3a induced NF κ B p65 expression in scrambled and *CaV1.2* shRNA (Fig. 5). On the other hand, *Tg737orp/ork* cells expressed a high basal level of NF κ B p65 which decreased in response to Wnt3a. In addition, *CaV1.2* expression was found to correlate with NF κ B p65.

CaV1.2 knockdown zebrafish develop PKD phenotypes

We have shown that *CaV1.2* localizes to primary cilia and have now elucidated the mechanism by which *CaV1.2* expression is regulated in renal epithelial cells. To assess the biological significance of *CaV1.2* expression, we used antisense

morpholinos to knockdown *CaV1.2* in zebrafish. Knockdown of the ciliary calcium channel polycystin-2 in zebrafish has been reported to result in PKD phenotypes including renal cyst formation, hydrocephalus, and left-right asymmetry (Obara et al., 2006). Our study showed that knockdown of *pkd2* increased *CaV1.2* expression (Fig. 6). This slight increase in *CaV1.2* was significant compared to the control morpholino. Interestingly, similar phenotypes were observed in *CaV1.2* morpholino (*cav1.2* MO) zebrafish. Compared with a non-specific control morpholino (control MO) injection, *cav1.2* MO zebrafish developed renal cysts (Fig. 7a), hydrocephalus (Fig. 7b) and various heart-looping defects (Fig. 7c). As generally accepted (Bakkers, 2011), left-right asymmetry was assessed by measuring the relative position of the cardiac atrium and ventricle with respect to the dorsal axis (Supplemental Movie 1).

Discussion

Non-motile primary cilia have been found to play a critical role in Wnt signaling by restricting β -catenin accumulation. Overexpression of polycystin-1 (a ciliary signaling receptor) inhibits GSK3 β and stabilizes β -catenin (Kim et al., 1999).




Fig. 4. Wnt3a increases ROS and DNA damage in *CaV1.2* shRNA but not in *Tg737orp/orpk* cells. **a:** Mitochondrial ROS was assessed by staining cells with MitoSOX. Wnt3a was found to increase ROS in scrambled and *CaV1.2* shRNA cells while decreasing ROS in *Tg737orp/orpk* cells when examined using fluorescence microscopy (bar = 20 μ m). **b:** Results were quantified using flow cytometry. **c:** Wnt3a increased formation of the oxidative DNA lesion 8-Oxoguanine in scrambled and *CaV1.2* shRNA cells but had no effect on *Tg737orp/orpk* cells (N = 3).

Polycystin-2 (encoded by *Pkd2*) is calcium channel forming protein found in primary cilia. In *Pkd2*^{-/-} embryos, cilia length was found to be decreased while β -catenin was upregulated (Kim et al., 2009). Interestingly, transgenic mice overexpressing β -catenin also developed cystic kidneys (Saadi-Kheddouci et al., 2001). Further, *LRP6*^{-/-} (a component of the Wnt receptor complex) mouse embryos die in utero with cystic kidneys (Pinson et al., 2000). Thus, primary cilia and Wnt signaling play a crucial role in PKD (Corbit et al., 2008).

Given that Wnt signaling also modulates mitochondrial physiology (Yoon et al., 2010), we examined the role of cilia in regards to mitochondria. *Tg737orp/orpk* contains an intron insertion at the 3' end of the intraflagellar transport 88 (*Ift88*) gene which results in a hypomorphic mutation that prevents ciliogenesis (Moyer et al., 1994). We used *Tg737orp/orpk* cells as a model for a cilia-deficient system. Through immunostaining, we confirmed the absence of cilia in *Tg737orp/orpk* cells compared with control (Fig. 1). The voltage-gated L-type calcium channel *CaV1.2* also localized to primary cilia. We generated a stable *CaV1.2* shRNA cell line and observed no changes in primary cilia compared with control. Further, treatment with Wnt3a had no effect on cilia number or length in scrambled or *CaV1.2* shRNA cells (data not shown). Thus, *CaV1.2* does not seem to play a role in ciliogenesis.

Mitochondrial biogenesis, oxidative phosphorylation, and generation of reactive oxidative species (ROS) were increased in response to Wnt3a in control renal epithelial cells

(Figures 2–4). The elevated levels of oxidative stress increased the formation of DNA lesions and the cellular DNA damage response (DDR). An interesting aspect of this response was an increase in expression of *CaV1.2* (Fig. 5). In *CaV1.2* knockdown cells, Wnt3a induced a similar effect on mitochondria and DDR. This data suggests that *CaV1.2* is a downstream effector in regard to Wnt signaling. In cilia-deficient cells, Wnt3a was unable to induce mitochondrial biogenesis and decreased mitochondrial activity, ROS production, and DDR. *CaV1.2* was found to be overexpressed in cilia-deficient cells as a compensatory mechanism; however, its expression decreased following Wnt3a treatment. Therefore, cilia length plays a role in regulating *CaV1.2* expression through modulation of Wnt signaling.

Defective primary cilia, indicated by either depletion of key ciliary proteins or fundamental changes in structure/length, results in PKD phenotypes (Wilson, 2004). Here we show that *CaV1.2* is a biologically significant ciliary protein. In the absence of *CaV1.2* in zebrafish (Fig. 6), PKD phenotypes including renal cyst formation, hydrocephalus, and left-right asymmetry defects were observed (Fig. 7). Moreover, *CaV1.2* was found to be overexpressed in *pkd2* knockdown zebrafish. This is intriguing given that both *CaV1.2* and PC2 are calcium channel forming proteins in the primary cilium, which further suggests a role for *CaV1.2* in PKD pathogenesis. Therefore, *CaV1.2* not only localizes to renal epithelial primary cilia, but it is also required for cilia function.




Fig. 5. Cilia modulates Wnt signaling to regulate CaV1.2 expression. Western blotting was performed on cellular protein extracts. Wnt3a treatment increased β-catenin accumulation in all cells, however, *Tg737^{orp/orp}* cells displayed high basal levels of β-catenin. NF-κB p65 was blotted as a readout for DNA damage response (DDR) to ROS induced DNA lesions. Wnt3a induced NF-κB p65 expression in scrambled and *CaV1.2* shRNA cells but decreased NF-κB p65 in *Tg737^{orp/orp}* cells. *CaV1.2* was overexpressed in *Tg737^{orp/orp}* cells. Wnt3a treatment induced CaV1.2 expression in scrambled shRNA while decreasing expression in *Tg737^{orp/orp}* cells. Data normalized to GAPDH for analysis (N = 3). Asterisks indicate significant difference from the corresponding control group ($P < 0.05$). # signs denote significant difference from the corresponding scrambled shRNA group ($P < 0.05$).




Fig. 6. *pkd2* knockdown increases CaV1.2 expression in zebrafish. Embryonic zebrafish protein extracts were obtained at 28 h postfertilization. Western blotting showed that *cav1.2* MO effectively reduced CaV1.2 expression compared with control MO. *pkd2* MO zebrafish were used to further verify morpholino specificity. Results were quantified through one-way ANOVA with Tukey post-test. Statistical significance is reported with a mean difference at the 0.05 level and denoted with an asterisk (N = 3).




a renal cyst**b hydrocephalus****c heart looping**

Fig. 7. CaV1.2 knockdown zebrafish develop renal cysts, hydrocephalus, and left-right asymmetry defects. CaV1.2 was knocked down in zebrafish (cav1.2 MO) using morpholino microinjection and compared to a scrambled control morpholino (control MO) to assess phenotypes. Renal cyst formation (a) and hydrocephalus (b) were measured at 3 days postfertilization through standard H&E staining as indicated by arrows. Left-right asymmetry was determined by measuring heart looping (c) at 2 days postfertilization under live microscopy.

Although it has been known that abnormal Wnt signaling leads to renal cyst formation (Lancaster et al., 2009), the underlying mechanism has been unclear. One proposed explanation is that Wnt signaling regulates renal cell proliferation and planar cell polarity to maintain renal tubule homeostasis (Happe et al., 2009). Our present study further suggests that cilia regulate Wnt signaling which ultimately controls CaV1.2 expression. Therefore, in the absence of CaV1.2, ciliary function is compromised leading to formation of renal cysts. In addition, our study also shows a previously unrecognized relationship between primary cilia and mitochondrial function. Overall, we propose that Wnt3a induces β -catenin accumulation in a cilia-dependent manner. This in turn increases mitochondrial biogenesis, oxidative phosphorylation, and generation of ROS that cause genomic DNA damage. The DDR triggers NF κ B p65 expression ultimately resulting in increased CaV1.2 expression.

Acknowledgments

This work was supported in part by DOD PR130153 and NIH DK080640.

Literature Cited

- Aberle H, Bauer A, Stappert J, Kispert A, Kemler R. 1997. Beta-catenin is a target for the ubiquitin-proteasome pathway. *EMBO J* 16:3797–3804.
- AbouAlaiwi WA, Takahashi M, Mell BR, Jones TJ, Ratnam S, Kolb RJ, Nauli SM. 2009. Ciliary polycystin-2 is a mechanosensitive calcium channel involved in nitric oxide signaling cascades. *Circ Res* 104:860–869.

- AbouAlaiwi WA, Muntean BS, Ratnam S, Joe B, Liu L, Booth RL, Rodriguez I, Herbert BS, Bacallao RL, Fruttiger M, Mak TW, Zhou J, Nauli SM. 2013. Survivin-induced abnormal ploidy contributes to cystic kidney and aneurysm formation. *Circulation*.
- Bai RK, Pong CL, Hsu CH, Wong LJ. 2004. Quantitative PCR analysis of mitochondrial DNA content in patients with mitochondrial disease. *Ann NY Acad Sci* 1011:304–309.
- Bakkers J. 2011. Zebrafish as a model to study cardiac development and human cardiac disease. *Cardiovasc Res* 91:279–288.
- Boveris A, Chance B. 1973. The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem J* 134:707–716.
- Boveris A, Oshino N, Chance B. 1972. The cellular production of hydrogen peroxide. *Biochem J* 128:617–630.
- Brown TA, Clayton DA. 2002. Release of replication termination controls mitochondrial DNA copy number after depletion with 2',3'-dideoxycytidine. *Nucleic acids Res* 30:2004–2010.
- Carter CS, Vogel TW, Zhang Q, Seo S, Swiderski RE, Moninger TO, Cassell MD, Thedens DR, Keppler-Noreuil KM, Nopoulos P, Nishimura DY, Searby CC, Bugge K, Sheffield VC. 2012. Abnormal development of NG2+PDGFR- α + neural progenitor cells leads to neonatal hydrocephalus in a ciliopathy mouse model. *Nat Med* 18:1797–1804.
- Corbit KC, Shyer AE, Dowdle WE, Gauden J, Singla V, Chen MH, Chuang PT, Reiter JF. 2008. Kif3a constrains beta-catenin-dependent Wnt signalling through dual ciliary and non-ciliary mechanisms. *Nat Cell Biol* 10:70–76.
- Gerdes JM, Liu Y, Zaghloul NA, Leitch CC, Lawson SS, Kato M, Beachy PA, Beales PL, DeMartino GN, Fisher S, Badano JL, Katsanis N. 2007. Disruption of the basal body compromises proteasomal function and perturbs intracellular Wnt response. *Nat Genet* 39:1350–1360.
- Happe H, Leonhard WN, van der Wal A, van de Water B, Lantinga-van Leeuwen IS, Breuning MH, de Heer E, Peters DJ. 2009. Toxic tubular injury in kidneys from Pkd1-deletion mice accelerates cystogenesis accompanied by dysregulated planar cell polarity and canonical Wnt signaling pathways. *Hum Mol Genet* 18:2532–2542.
- Jackson SP, Bartek J. 2009. The DNA-damage response in human biology and disease. *Nature* 461:1071–1078.
- Janssens S, Tschopp J. 2006. Signals from within: The DNA-damage-induced NF- κ B response. *Cell Death Differ* 13:773–784.
- Jin X, Mohieldin AM, Muntean BS, Green JA, Shah JV, Mykityn K, Nauli SM. 2013. Cilioplasm is a cellular compartment for calcium signaling in response to mechanical and chemical stimuli. *Cell Mol Life Sci*.
- Kasai H. 1997. Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis. *Mutat Res* 387:147–163.
- Kasai H, Tanooka H, Nishimura S. 1984. Formation of 8-hydroxyguanine residues in DNA by X-irradiation. *Gann* 75:1037–1039.

- Kim E, Arnould T, Sellin LK, Benzing T, Fan MJ, Gruning W, Sokol SY, Drummond I, Walz G. 1999. The polycystic kidney disease 1 gene product modulates Wnt signaling. *J Biol Chem* 274:4947–4953.
- Kim I, Ding T, Fu Y, Li C, Cui L, Li A, Lian P, Liang D, Wang DW, Guo C, Ma J, Zhao P, Coffey RJ, Zhan Q, Wu G. 2009. Conditional mutation of Pkd2 causes cystogenesis and upregulates beta-catenin. *J Am Soc Nephrol* 20:2556–2569.
- Lancaster MA, Louie CM, Silhavy JL, Sintasath L, Decambre M, Nigam SK, Willert K, Gleeson JG. 2009. Impaired Wnt-beta-catenin signaling disrupts adult renal homeostasis and leads to cystic kidney ciliopathy. *Nat Med* 15:1046–1054.
- Lancaster MA, Schroth J, Gleeson JG. 2011. Subcellular spatial regulation of canonical Wnt signalling at the primary cilium. *Nat Cell Biol* 13:700–707.
- Moyer JH, Lee-Tischler MJ, Kwon HY, Schrick JJ, Avner ED, Sweeney WE, Godfrey VL, Cacheiro NL, Wilkinson JE, Woychik RP. 1994. Candidate gene associated with a mutation causing recessive polycystic kidney disease in mice. *Science* 264:1329–1333.
- Muntean BS, Horvat CM, Behler JH, Aboulaiwi WA, Nauli AM, Williams FE, Nauli SM. 2010. A comparative study of embedded and anesthetized zebrafish in vivo on myocardial calcium oscillation and heart muscle contraction. *Front Pharmacol* 1:139.
- Muntean BS, Jin X, Nauli SM. Chapter 9: Primary cilia are mechanosensory organelles with chemosensory roles. *Mechanosensitivity and mechanotransduction*. ISBN: 978-994-007-2003-2009.
- Nauli SM, Alenghat FJ, Luo Y, Williams E, Vassilev P, Li X, Elia AE, Lu W, Brown EM, Quinn SJ, Ingber DE, Zhou J. 2003. Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet* 33:129–137.
- Nauli SM, Kawanabe Y, Kaminski JJ, Pearce WJ, Ingber DE, Zhou J. 2008. Endothelial cilia are fluid shear sensors that regulate calcium signaling and nitric oxide production through polycystin-1. *Circulation* 117:1161–1171.
- Norris DP. 2012. Cilia, calcium and the basis of left-right asymmetry. *BMC Biol* 10:102.
- Obara T, Mangos S, Liu Y, Zhao J, Wiessner S, Kramer-Zucker AG, Olale F, Schier AF, Drummond IA. 2006. Polycystin-2 immunolocalization and function in zebrafish. *J Am Soc Nephrol* 17:2706–2718.
- Pendergrass W, Wolf N, Poot M. 2004. Efficacy of MitoTracker Green and CMXRosamine to measure changes in mitochondrial membrane potentials in living cells and tissues. *Cytometry A* 61:162–169.
- Pinson KI, Brennan J, Monkley S, Avery BJ, Skarnes WC. 2000. An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature* 407:535–538.
- Poot M, Pierce RH. 2001. Analysis of mitochondria by flow cytometry. *Methods Cell Biol* 64:117–128.
- Saadi-Kheddouci S, Berrebi D, Romagnolo B, Cluzeaud F, Peuchmaur M, Kahn A, Vandewalle A, Perret C. 2001. Early development of polycystic kidney disease in transgenic mice expressing an activated mutant of the beta-catenin gene. *Oncogene* 20:5972–5981.
- Wilson PD. 2004. Polycystic kidney disease. *N Engl J Med* 350:151–164.
- Yoon JC, Ng A, Kim BH, Bianco A, Xavier RJ, Elledge SJ. 2010. Wnt signaling regulates mitochondrial physiology and insulin sensitivity. *Genes Dev* 24:1507–1518.

Supporting Information


Additional supporting information may be found in the online version of this article at the publisher's web-site.

Vascular Endothelial Primary Cilia: Mechanosensation and Hypertension

Ashraf M. Mohieldin^{2,#}, Hossain Saad Md Zubayer^{1,#}, Alzahra J. Al Omran¹, Hannah C. Saternos¹, Ali Zarban¹, Surya M. Nauli³ and Wissam A. AbouAlaiwi^{1,*}

¹Department of Pharmacology, University of Toledo Health Science Campus, Toledo, OH, USA;

²Department of Medicinal & Biological Chemistry, University of Toledo Health Science Campus, Toledo, OH, USA; ³Department of Biomedical & Pharmaceutical Sciences, Chapman University Rinker Health Science campus, Irvine, CA, USA



W.A. AbouAlaiwi

Abstract: Primary cilia are sensory organelles that extend from the cell surface and sense extracellular signals. Endothelial primary cilia protruding from the inner surface of blood vessel walls sense changes in blood flow and convert this mechanosensation into an intracellular biochemical/molecular signal, which triggers a cellular response. Primary endothelial cilia dysfunction may contribute to the impairment of this response and thus be directly implicated in the development of vascular abnormalities such as hypertension and aneurysms. Using both *in vitro* techniques as well as *in vivo* animal models, we and others have investigated fluid flow mechanosensory functions of endothelial cilia in cultured cells, animal models and autosomal dominant polycystic kidney disease (ADPKD) patients. More in-depth studies directed at identification of the mechanisms of fluid flow sensing will further enhance our knowledge of cilia-dependent vascular pathology. Although the current treatments aimed at treating the cardiovascular symptoms in ADPKD patients successfully slowed the progression of cyst growth, there is growing evidence which suggests that drugs which interfere with primary cilia function or structure could reduce cardiovascular complications in ADPKD. This review is to summarize the most recent studies on primary endothelial cilia function in the vascular system and to present primary cilia as a novel therapeutic target for vascular hypertension.

Keywords: Cardiovascular, cell division, fluid-shear, hypertension, primary cilia.

INTRODUCTION

It has been over a century since primary cilia have been visualized, but the study of their sensory role is, comparatively, a new field [1]. Like other cellular organelles i.e. nucleus, Golgi bodies, mitochondria etc. cilia can be considered as a separate entity and have the following specialized functions:

- Mechanosensation [2-4].
- Shear-stress sensation [5-7].
- Nodal flow generation [8-10].
- Nodal flow sensation [11, 12].
- Osmosensation [13-15].
- Chemosensation [16-19].
- Light sensation [20-22].
- Smell sensation [23, 24].
- Gravitational sensation [25].
- Developmental regulation [26-28].
- Fluid clearance [29, 30].
- Sperm motility [31-33].
- Fluid transportation [34-36].
- Oocyte transportation [37, 38].
- Sound-wave sensation [39-41].

In order to study a cilium, it must be divided into five distinct structural blocks – the ciliary membrane, the soluble compartment, the axoneme, the ciliary tip and the basal body complex (Fig. 1a). The primary ciliary membrane has a distinct lipid bilayer composition and is continuous with the cell membrane at the transition fiber. It houses various membrane mechanosensory receptors, ligand-activated chemoreceptors and ion channels to support mechanosensation, chemosensation and ion influx in response to extracellular stimuli. The soluble or matrix compartment also called the cilioplasm is composed of fluid materials to support signaling activities. The axoneme, which emanates from the basal body, is composed of nine pairs of microtubules forming heterodimer structure. Apart from delivering cellular components in and out of the ciliary shaft, it plays a significant role in maintaining a long ciliary structure. The ciliary tip contains specialized proteins, but

*Address correspondence to this author at the Department of Pharmacology; MS 1015, The University of Toledo, College of Pharmacy and Pharmaceutical Sciences, Health Education building; Room 282E, 3000 Arlington Ave, Toledo, OH 43614; Tel: 419-383-1949; Fax: 419-383-1909; E-mail: Wissam.Abou-Alaiwi@UToledo.Edu

[#]Hossain Saad Md Zubayer and Ashraf M. Mohieldin contributed equally to this work.




Fig. (1). Mechanosensory cilium. A primary cilium is a cellular organelle extending from the apical surface of many cell types. The mechanosensory cilia act as antennas to receive and transmit extracellular fluid-shear stress into cellular biochemical signals. **(a)** A cilium is composed of a ciliary membrane, a soluble matrix, a ciliary axoneme, and a basal body complex. The *ciliary membrane* contains various sensory proteins. The *soluble matrix* contains many regulatory proteins that are involved in signal transduction networks. The *ciliary axoneme* is surrounded by the nine pairs of microtubules that are thought to control the length of a cilium. The *basal body complex* is the anchoring point for axonemal microtubules. **(b)** Primary cilia found on the surface of endothelial cells can be localized by a simple immunostaining experiment using antibody against acetylated α -tubulin (green) to label primary cilia and pericentrin (red) to label centriole or basal body. The nucleus is counterstained with DAPI to label DNA. **(c)** Mouse primary epithelial cilia can be studied *in vivo* by scanning electron microscopy (SEM) to reveal kidney epithelial cilia protruding from the surface of kidney epithelial cells towards the lumen of kidney tubules. **(d)** Primary cilia are sensory organelles that sense fluid-shear stress on the apical surface of cells. Fluid flow that produces enough drag-force on the cells will bend sensory cilia. Part of this figure is reproduced from [152].

their roles are still unknown. The basal body complex is composed of the mother and daughter centrioles connected by rootlets and emanate from the basal body anchorage. The mother centriole is the basal body from which the cilium protrudes [42].

Primary Cilia is a term commonly used to denote “9+0” non-motile cilia. These have nine parallel doublet microtubules (denoted by ‘9’) and a central pair of microtubule is absent within the central sheath (denoted by ‘0’). Length of a primary cilium can vary from 2-50 μm and it has a diameter of approximately 0.25 μm [42]. Primary cilia are found on several mammalian cell types i.e. endothelia (Fig. 1b) [43-46], epithelia (Fig. 1c) [47-49], neurons [50-53], osteoblasts [54-56], fibroblasts and others [57-59]. Thus, abnormal ciliary proteins and/or abnormal ciliary function can be associated with various pathological conditions collectively known as ciliopathies; which include polycystic kidney disease (PKD), nephronophthisis, bardet biedl syndrome, left-right asymmetry defect, oral facial syndrome, obesity, hypertension and others [60]. Primary cilia can be activated by bending through perfusing cells with fluid-shear stress (Fig. 1d) or by mechanical stress [61]. Bending cilia by apical fluid perfusion with physiologically relevant flow rate [2, 62] or by suction through a micropipette [63] can activate primary cilia. Consequently, mechanosensory ciliary proteins such as polycystin-1 [5], polycystin-2 [7], transient receptor potential-4 [64], fibrocystin [65], and probably many others convert this mechanical force into an intracellular biochemical signal [66, 67].

PRIMARY ENDOTHELIAL CILIA AS BLOOD PRESSURE SENSORS

The control of the circulatory system depends largely on the mechanical properties of blood vessels in both normal

and pathological conditions. The continuous change in blood vessel diameter triggered by contraction and relaxation of vascular smooth muscle cells is important for normal blood flow [68-71].

Different blood vessels are subjected to changes in blood pressure as well as changes in the hemodynamic forces which induce distinct responses within the vascular tree and allows for tissue perfusion. Resistant arteries have the most influence on blood flow and they are subjected to the continuous effect of blood flow and pressure. An increase in blood pressure induces a vasoconstriction known as the myogenic tone, which facilitates the action of the sympathetic nervous system [72]. When transmural blood pressure increases, vessel diameter is reduced [73-75]; while faster flow velocity (shear stress) increases blood vessel diameter [74, 76, 77]. Vascular endothelial cells lining the inner wall of blood vessels can sense changes in blood flow and pressure and convert these mechanical changes into changes of smooth muscle tone [74, 78].

The pulsatile nature of blood flow through the vasculature exerts different types of mechanical forces such as shear stress, cyclic strains and hydrostatic pressure that can impact the physiology of the blood vessel wall. Being the inner most layer and in direct contact with blood flow, the endothelial cell layer bears the most frictional forces induced by the flow of blood. Though these forces are practically impossible to differentiate *in vivo*, they can be distinguished from one another in *in vitro* and *ex vivo* studies [79-81]. In this review, we will focus on shear stress which is the most studied mechanical force in cilia research.

We have previously demonstrated the presence as well as the function of endothelial primary cilia as fluid mechanosensor *in vitro* in mouse aortic endothelial cells, *ex*

in vivo in isolated mouse arteries and *in vivo* in mouse models as well as in blood vessels from human patients [5, 7]. The presence as well as the size of primary endothelial cilia varies with respect to the level of fluid shear stress or fluid turbulence. For example, blood vessels with relatively low fluid shear stress have longer cilia while blood vessels with high fluid shear stress are devoid of cilia or have shorter cilia simply due to the inability of cilia to tolerate high levels of shear stress, which could induce its disassembly. Consequently, the ciliary mechanosensory function in those areas of high fluid shear stress could be minimized or substituted by other vessel components such as the glycocalyx [82]. Sensory proteins localized to primary endothelial cilia such as polycystins can detect any sudden increase in blood pressure or shear stress inside the blood vessel [5, 7]. This in turn, leads to the influx of calcium ions to the cilium followed by an increase in intracellular calcium concentration. As a result, primary cilia enable cells to translate this extracellular fluid mechanics into a complex intracellular signaling cascade. This signaling cascade eventually leads to the activation of endothelial nitric oxide synthase (eNOS), an endothelial enzyme that synthesizes nitric oxide gas. Nitric oxide diffuses from endothelial cells to the surrounding smooth muscle cells and promotes vasodilation [83-85]. Both polycystin-1 and polycystin-2, an 11 transmembrane protein with a long extracellular domain and a cation channel with 6 transmembrane domains, respectively are expressed in the endothelial and vascular smooth muscle cells of all major blood vessels [86]. We previously reported the loss of response to fluid-shear stress in mouse endothelial cells with *knockdown* or *knockout* of *PKD2* [7]. In addition to the mouse data, polycystin-2 null endothelial cells generated from *PKD2* patients that do not show polycystin-2 in the cilia are unable to sense fluid flow. Therefore, mutations in both *PKD1* and *PKD2* have been shown to contribute to hypertension [87], in part by the failure to convert an increase in mechanical blood flow into cellular NO biosynthesis [5, 7]. In summary, by sensing any increase in fluid shear stress, primary cilia can activate different cellular mechanisms in order to lower total peripheral resistance and consequently, blood pressure.

It is crucial to discern the physiological mechanism behind mechanosensory role of primary cilia in order to clearly understand the correlation between primary cilia abnormalities and cardiovascular pathology and in particular hypertension. A new mechanism has been proposed that involves polycystin-1 and polycystin-2 [5, 7]. Endothelial cells without primary cilia were isolated from *Tg737^{Orpk/Orpk}* mice to examine the mechanosensory role of primary cilia. In addition and to confirm the role of polycystin-1 and polycystin-2 as mechanosensory proteins, endothelial cells were collected from PKD mice and ADPKD human patients. Polycystins are absent in the cilia of these cells and present in *Tg737* mice cells, but localized at the base of the “stubby” primary cilia. Different magnitudes of fluid shear stress were applied to these cells. Though shear induced cytosolic calcium was increased in normal endothelial cells, diseased artery and mutant endothelial cells did not show any calcium response to fluid shear stress [5, 7]. To verify cilia function specifically against fluid shear stress, a new technique involving artery perfusion in glass capillary was utilized [7,

88]. It was observed that though cilia react normally when other mechanical stimuli are applied, response to fluid shear stress is highly altered.

To dissect the molecular mechanisms downstream of cilia function, different biochemical and pharmacological inhibitors were used in these studies to block downstream molecular targets in endothelial cells. When EGTA is used to block extracellular calcium, both cytosolic calcium and nitric oxide production was inhibited. In addition, L-NAME (N^G -nitro-L-arginine methylester), an eNOS inhibitor, was found to block shear induced nitric oxide synthesis but not cytosolic calcium increase. This indicates the necessity of extracellular calcium influx for the downstream generation of nitric oxide. To examine calcium dependent mechanism of nitric oxide biosynthesis, calphostin C, a PKC inhibitor and W7, a calmodulin antagonist were employed. Results showed that PKC and calmodulin function downstream of calcium pathway and blocking either of these will inhibit nitric oxide synthesis. When pharmacological blockers Akt inhibitor II, LY-294,002, and wortmannin were applied, role of Akt was confirmed in shear induced nitric oxide production. These studies in addition to others demonstrate an interaction between mechanosensory polycystin-1 and polycystin-2 and that absence of this interaction results in disturbance of localization of polycystin-2 to primary cilia and hence, disturb ciliary functions. In summary, these studies lead us to propose that, after sensing fluid shear stress, polycystin-1 interacts with and activates polycystin-2, a calcium channel which induces an intracellular calcium influx. This is followed by activation of PKC and formation of calcium-calmodulin complex. Our data also indicate that Akt/PKB is also involved in regulation of eNOS activation and the generation of nitric oxide in response to fluid shear (Fig. 2a and b).

ROLE OF PRIMARY CILIA IN THE CARDIOVASCULAR SYSTEM

Despite its well-known and demonstrated function as a mechanosensory organelle in the cardiovascular system, the role of primary cilia in this system is still a widely debated topic. Recent studies have successfully demonstrated the presence of primary cilia throughout the cardiovascular system. Primary cilia have been observed in endocardia [6, 89], arterial endothelia [2, 7], venous endothelia [90], corneal endothelia [44, 46], arterial smooth muscle cells [91] and airway smooth muscle cells [92].

Primary Cilia and Heart Development

Recent studies have demonstrated an important role of cardiac primary cilia in harmonizing heart development, and defects in these cilia are associated with inherited heart disease. Different types of cilia are present in different chambers throughout heart development to control cardiogenesis. For example, sensory nodal cilia play an important role in regulating signaling mechanisms essential for the establishment of left-right asymmetry, a process for regulating heart morphogenesis. In addition, primary cilia are present on the surface of cardiomyocytes during heart development and houses different receptor complexes which coordinate various signaling mechanisms that are important




Fig. (2). Endothelial primary cilia as regulators of blood pressure through nitric oxide production and vascular architecture through survivin. (a and b) Nitric oxide synthesis and survivin expression depend on the presence of functional sensory cilia housing sensory complex, polycystin-1 and polycystin-2. Bending of primary endothelial cilia by fluid flow activates polycystin complex and initiates a series of intracellular signaling cascades resulting in the synthesis and release of NO gas, which vasodilates smooth muscles in the vasculature. These intracellular signaling cascades include calcium influx, calcium-calmodulin, protein kinase C, Akt/PKB and eNOS. Survivin downregulation is involved in aneurysm formation. Fluid shear stress induced the activation of polycystin-1/2 complex followed by increase in intracellular calcium influx. Akt and NF-κB are downstream effectors of PKC pathway in regulating flow-induced survivin expression. The increase in p-NF-κB expression is accompanied by an increase in survivin expression, which is critical for normal endothelial cell division and consequently vascular integrity. Modified from [152, 153] with permission.

for proper heart morphogenesis and development. Cilia mutants have been characterized by heart developmental defects such as ventricular and atrial septum defects, myocardial wall disorganization and thinning and double-outlet right ventricles due to mutations in *pkd1* or *pkd2*, the causative genes of ADPKD [93]. Moreover, knockout mice of cilia structural genes such as *ift88* and *kif3a* are also characterized by severe cardiac phenotypes such as hypoplasia of the endocardial cushions, reduction in ventricular trabeculation and enlarged pericardium [94]. This suggests that these cardiac phenotypes could be contributed mainly by ciliary dysfunction and that primary cardiac cilia are critical for normal heart development. More importantly, dysfunctional cilia are associated with congenital heart disease. Several ciliopathies such as nephronophthisis (NPHP), Meckel syndrome (MKS), Bardet-Biedl syndrome (BBS), and Joubert syndrome (JBTS) are usually associated with cardiac defects such as septum defects and aortic stenosis [95].

Cardiac development is controlled by complex regulatory networks of both inductive and inhibitory signals from both the heart as well as the surrounding tissue which coordinate different stages of heart development. An example is the Hedgehog (Hh) signaling pathway which regulates the L-R asymmetry in vertebrates and promotes the activation of several transcription factors involved in various cellular mechanisms during tissue homeostasis and development. It

has been shown that primary cilia can regulate the Hh signaling pathway in several cell types by functioning as a unique compartment for the continuous turnover of Hh signaling components. As a result, defects in primary cilia assembly and consequently Hh signaling components turnover can lead to a variety of developmental pathologies including congenital heart defects [96]. Another example of a signaling network which plays a critical role during heart development is the TGFβ/Bone Morphogenic Protein (BMP) network. It has been recently shown that primary cilia can regulate the canonical TGFβ signaling network through the activation of transcription factors Smad2/3 [97]. Furthermore, it has been shown that the TGFβ ligand, TGFβ-1 can stimulate the differentiation of stem cells into cardiomyocytes and that *Tg737^{Orpk/Orpk}* mouse embryonic fibroblasts are characterized by diminished TGFβ activity, suggesting a direct role of cardiac primary cilia in regulating TGFβ signaling during cardiomyogenesis. Taken together, these studies prompted us to speculate that primary cilia may function as signaling hubs facilitating the cross-talk between the different signaling networks that regulate heart development.

Primary Endothelial Cilia and Vascular Architecture

The morphology and function of blood vessels are continuously altered in response to blood flow which is mainly detected by the vascular endothelium. An aneurysm

is a balloon-like bulge in the wall of blood vessels. Studies on both ADPKD patients and animal models indicate that polycystins are essential for normal blood vessel's structure and development [98]. Aneurysm formation is one of the deadliest vascular abnormalities in PKD patients. The occurrence of aneurysm represents a major risk factor for morbidity and mortality associated with PKD [99]. Mice with *Pkd* mutations develop hemorrhages and aneurysms [100]. Vascular aneurysms are associated with tissue remodeling due to abnormal proliferation of the endothelial cell layers in response to complex hemodynamic changes in fluid shear stress [101]. We recently demonstrated that aneurysm formation is more severe in *survivin*, *pkd1* and *Tg737* vascular-specific knockout (*Pdgfr β Cre:Survivin^{flox/flox}*, *Pdgfr β Cre:Pkdl^{flox/flox}* and *Pdgfr β Cre:Tg737^{flox/flox}*) mice than in wild type mice [102]. An abnormality of primary cilia structure (polaris) or function (polycystin-1) is associated with downregulation of survivin expression, which is associated with abnormal mitotic events in endothelial cells [103]. This in turn can lead to abnormal cytokinesis triggered by polyploidy formation and ultimately contributes to vascular aneurysm [102]. Thus, the inability of primary endothelial cilia to respond to fluid shear stress can contribute to the exacerbation of aneurysm formation in these mice models.

Protein Kinase C (PKC) and Akt are downstream signaling messengers of primary cilia. Studies on endothelial cells have verified that survivin expression following cilia activation is regulated by PKC, Akt and Nuclear Factor- κ B (NF- κ B). Akt is downstream of PKC and can regulate NF- κ B, which is known to regulate survivin expression. Taken all the above together, we propose a Cilia-PKC-Akt-NF- κ B pathway involved in survivin expression and cell division regulation [102, 104-107]. All in all, the inability of mechanosensory cilia to detect fluid shear stress is associated with down-regulation of survivin leading to abnormal cytokinesis triggered by polyploidy formation and finally contributing to vascular aneurysm (Fig. 2b) [102].

Primary Cilia in Corneal Endothelium

The corneal endothelium located beneath the anterior chamber of the eye is orchestrated into a two-dimensional layer of hexagonal cells and plays critical role in maintaining corneal transparency. The presence of primary cilia in the corneal endothelium has been demonstrated in various mammalian species [44]. In humans, corneal endothelial cilia protruding into the anterior chamber of the eye have similar structure to the "9+0" primary cilia, but their function remains somehow unclear, although evidence for an osmoregulatory or a chemosensory function exists. Moreover, it has been postulated that primary cilia of the mouse corneal endothelium play key role towards proper morphogenesis of the characteristic hexagonal shape or pattern of the corneal endothelium during development and disassemble following tissue homeostasis during adult stage pointing towards a role of corneal endothelial primary cilium in corneal endothelial development [108].

More importantly, a link between primary cilia and intraocular pressure regulation has been recently demonstrated through studying a rare X-linked congenital

syndrome known as pediatric glaucoma or Lowe syndrome [109]. It has been previously shown that the *Oculocerebrorenal syndrome of Lowe (OCRL)* gene encoding an inositol polyphosphate 5- phosphatase, which is mutated in Lowe syndrome is involved in vesicle trafficking to the primary cilium [110]. More recently, a potential role of primary cilia in the trabecular meshwork which regulates intraocular pressure in the eye has been demonstrated. In this study, it was shown that primary cilia are important sensors of intraocular pressure changes and that in glaucoma, defective trabecular meshwork cells have shortened primary cilia due to increased intraocular pressure and fluid flow [109]. Consequently, this shortening is associated with increased expression of TNF α , TGF β and GLI1 genes. Primary cilia response to increased intraocular pressure is mainly mediated through interaction between OCRL and transient receptor potential vanilloid 4 (TRPV4), a ciliary mechanosensory channel. Although these findings significantly advance the current understanding of how intraocular pressure is regulated, they provide further evidence and support for our proposed role of primary endothelial cilia as both blood pressure sensor and regulator.

Primary Cilia and Vascular Smooth Muscle Cell Function

Vascular smooth muscle cells (VSMCs) localized in the medial layer of the arterial wall play critical roles in maintaining and regulating blood vessel tone, blood pressure and blood flow [111]. VSMCs express a unique group of proteins with specialized contractile functions and have very low proliferative profile during adult stages. The environment in which VSMCs reside is mainly composed of collagen and elastic fibers as extracellular matrix (ECM) components. Normally, VSMCs are not exposed directly to fluid shear stress simply because the inner lining endothelial cell layer provide the contacting surface with blood flow and consequently shields the VSMC layer. However, in cases of endothelial cell layer injury or denudation, VSMCs become directly exposed to fluid shear stress levels similar to the endothelial cell layer in intact vascular regions. It is critical to understand the mechanisms by which VSMCs sense and transduce the stimuli of shear stress into intracellular biochemical signals which allows better understanding of vascular disease. Studies reviewing the effects of fluid shear stress on VSMCs revealed the presence of shear specific mechanosensors such as cell membrane-related receptors, cell surface glycocalyx, ion channels and integrins as well as the presence of shear-specific secondary signaling messengers such as NO, Ca⁺² and MAPKs. These messengers in turn regulate the expression of shear-responsive genes which control various VSMCs responses such as proliferation, differentiation, apoptosis and migration.

A sensory role of primary cilia in VSMCs similar to its role in several other cellular systems has not been demonstrated until recently [112]. It has been shown that primary cilia are localized to VSMCs in a preferentially oriented pattern with respect to the artery axis as well as to the ECM and they house mechanosensory proteins that are responsive to ECM proteins and to cell-ECM interaction. These initial observations suggest a mechanochemical

sensory role of VSMCs primary cilia in the vasculature. In addition, primary cilia on the surface of VSMCs may act as fluid mechanosensors regulating intracellular Ca^{+2} influx and VSMCs migration in response to fluid shear stress. Primary cilia and cell surface glycocalyx are both displaced in response to fluid flow and affect cell surface receptors as well as integrins. Both cell contraction and migration requires disassembly of integrin-mediated focal adhesion [113]. Therefore, it remains to be discovered if VSMCs primary cilia can sense interstitial flow and play a role in controlling VSMCs contraction and migration. In addition, VSMCs *in vivo* are exposed to other hemodynamic forces such as stretch, and pressure at the same time, which may act in concert to regulate mechanosensitive signaling pathways controlling vascular function and disease. In summary, it is safe to assume that if VSMCs' primary cilia function or structure is perturbed, it will contribute to vascular disease and hypertension.

HYPERTENSION AND HUMAN ADPKD

ADPKD is an example of a ciliopathy or a genetic disorder resulting from the abnormal function and/or structure of primary cilia. Even with successful renal transplant or replacement therapy, patients with ADPKD eventually die due to cardiovascular complications, including hypertension, aneurysms, aortic root dilation, dissection, vascular ectasia, mitral valve prolapse and abnormal function of the microvascular bed [99, 114-116], heart failure and congestive heart disease [115-121]. Aside from the kidney phenotype, the frequencies of phenotypic manifestations in patients with ciliopathy include congestive heart failure/hypertension (78%), hepatic cysts (75%), diverticulosis (70%), ovarian cysts (40%), cardiac valve disorders (25%), inguinal hernias (15%) and intracranial aneurysms (10%) [107]. Our group is one of the first to report cilia function using mouse models and patient samples to explain cystic kidney phenotypes [2, 122] and to report cilia function in nitric oxide (NO) biosynthesis [5, 123]. Plasma concentration of the vasodilator nitric oxide has been found to be lowered in ADPKD patients compared to healthy volunteers [124]. This demonstrates a strong link between endothelial dysfunction and ADPKD. Another study reported similar results where the level of ADMA (asymmetric dimethylarginine), a marker of an inhibitor of nitric oxide synthase, was found to be elevated in PKD patients [125].

The correlation between hypertension and kidney volume occurs in the early childhood stages of ADPKD. The development of hypertension in ADPKD patients is associated with increased renal volume as well as increased left ventricular mass index. Similarly, it has also been suggested that high blood pressure can advance cyst growth [126, 127]. Interestingly, hypertension appears in children before they confront any significant reduction in glomerular filtration rate or development of ADPKD [128, 129]. Now that ADPKD is strongly correlated with dysfunction of ciliary proteins and/or abnormal cilia structure [2, 130, 131], this prompted us to propose that, endothelial dysfunction in ADPKD patients may be attributed to the inability of

primary endothelial cilia to sense fluid-shear stress, or to an anomaly in any other downstream signaling mechanism. Taken together, this suggests that the pathogenesis of hypertension is a risk factor of endothelial primary cilia dysfunction at least partially independent from kidney function in ADPKD.

On the other hand, renal cyst enlargement in ADPKD adults has been ascribed to contribute in part to hypertension. Enlargement of renal cysts leads to the stimulation of both circulating as well as the intrarenal renin-angiotensin-aldosterone system (RAAS) system due to the compression of the cyst-adjacent parenchyma on the renal vasculature resulting in areas of renal ischemia and vascular changes [132, 133]. Activation of RAAS in PKD has been substantiated in both clinical studies [134] and murine models [135, 136]. Other components of the RAAS, including angiotensinogen, angiotensin converting enzyme (ACE), angiotensin II receptor and angiotensin II peptide have also been detected in cysts and dilated tubules in ADPKD kidneys [137]. In addition to hypertension, angiotensin contributes significantly to cyst growth and expansion, increased endothelin levels, fibrosis and increased sympathetic activity. Sympathetic activity can stimulate RAAS and angiotensin can stimulate sympathetic nervous system as well [138, 139]. Moreover, activation of the RAAS has been found in normotensive and hypertensive PKD patients, regardless of their blood pressure and renal function [140]. Not surprisingly, the high levels of circulating angiotensin II in PKD patients have been shown to contribute to the development of vascular hypertrophy, which is further implicated in vascular remodeling [141]. The effect of blockage of the renin-angiotensin-aldosterone system with Angiotensin Converting Enzyme inhibitors (ACE-I) and angiotensin receptor antagonists on renal volume and kidney function is discussed in the next section. A study involving eleven kidney transplant cases of hypertensive PKD patients accounted an improved blood pressure in only six patients [142]. Another report demonstrated that PKD patients with hypertension continue to show cardiovascular complications after renal transplantation [143]. These studies clearly suggest that, though kidney transplantation is beneficial for improving some cardiovascular issues, it is not sufficient to control hypertension in PKD patients. Whether hypertension contributes to diminished renal function or deteriorated kidney function exacerbates hypertension is still a debated topic that warrants further discussion and there is a wide scope of more research to clearly understand the cause and effect phenomenon undergoing in this case.

AVAILABLE AND ONGOING TREATMENT OPTIONS FOR CARDIOVASCULAR DISEASES IN ADPKD

A recent review involving 1877 ADPKD patients' records showed that the use of antihypertensive therapy in ADPKD patients have been increased from 32% in 1991 to 62% in 2008. A similar increase in the use of RAAS inhibitors was also reported (from 7 % to 42 %) [144]. But given that cardiovascular complications are the most

common cause of death in ADPKD patients, it is justified that researches are going on to a greater extent for innovation of novel therapeutics.

Alike regular hypertensive patients, reduced salt and caffeine intake, smoking cessation, regular exercise, and optimal water consumption have been suggested and found beneficial in hypertensive PKD patients [145, 146]. Pathophysiology of hypertension in this disease suggests the usefulness of ACE inhibitors. ACE-I enalapril is found to control blood pressure but it has a consistent antiproteinuric effect [133]. In a different study, 3% of patients displayed a reversible increase in serum creatinine after initiation of ACE-I use [147, 148]. So, ACE-Is should be used cautiously in ADPKD patients. The RAAS based pathophysiology of hypertension foretells the effectiveness of Angiotensin Receptor Blockers (ARBs). A randomized trial compared candesartan (ARB) with amlodipine (calcium channel blocker) and both were found equally effective in controlling blood pressure in PKD patients [149]. However, there is no other study exhibiting similar outcome and should be explored further. Dual blockade of RAAS with ACE-Is and ARBs has also been suggested and being tested in two clinical trials [150]. The most recent release of the results of the HALT study among 558 hypertensive participants with ADPKD revealed that the combination of lisinopril, an ACE-I and telmisartan, an ARB did not significantly alter the rate of increase in total kidney volume in the early stages of ADPKD. However, they indicated that aggressive blood-pressure control might slow kidney cyst growth and reduce left-ventricular-mass index compared to normal blood pressure control [151]. Beta-blockers (metoprolol, atenolol) and calcium channel blockers (amlodipine) are found to be effective in controlling hypertension in PKD patients. Though metoprolol did not alter glomerular filtration rate (GFR) or urinary albumin excretion, effects of atenolol on kidney function is still unknown. Calcium channel blockers are not recommended for use as a first line therapy in PKD patients, as this may cause a significant reduction in GFR. Theoretically, diuretics are recommended in PKD patients only as an additional therapy with ACE-Is as diuretics activate RAAS. Moreover, patients treated with diuretics showed increased urinary excretion of proteins [150].

We and others have independently studied abnormality in primary cilia in the vascular and renal systems in PKD. We hypothesized that cilia in the vascular system play important roles in regulating blood pressure and vascular integrity through the expression of mechano- or chemosensory receptors. Supporting our hypothesis is that patients with abnormal cilia function (PKD) also show a greater incidence of hypertension and other cardiovascular problems. In theory, drugs that interfere or alter cilia sensory function could prevent or delay this high blood pressure increase. This idea is based on our most recent chemical-screening studies showing that dopamine receptor is localized in endothelial cilia and that this ciliary receptor has dual chemo and mechano-sensing roles in endothelial cells. We showed that “activated” dopamine receptor-type 5 in the cilia could switch its chemosensing function to mechanoreceptor, while “non-activated” receptor has a primary function as a chemoreceptor. Activation of ciliary dopamine receptor

could bypass mechano-ciliary polycystin complex through a mechanism that increases the cilia length. The extension of cilia probably serves to increase the sensitivity of these sensory organelles to fluid flow, in which dopamine receptor-type 5 acts as the mechanoreceptor.

CONCLUSION

As more evidence is presented to support the direct involvement of primary cilia in the cardiovascular system, this necessitates large-scale screening studies for drugs to investigate whether targeting primary cilia function or structure towards hypertension results in better outcomes.

CONFLICT OF INTEREST

The author(s) confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Authors would like to thank Charisse Montgomery for her editing service. Due to restricted space, the authors apologize to those whose work is not cited in this paper. Z. Saad's and A. Mohieldin's work partially fulfills the requirements for a Ph.D. degree in Experimental Therapeutics and Medicinal and Biological Chemistry degrees, respectively. A. Al-Omran's, A. Zarban's and H. Saternos's work fulfills the requirements for a Master's degree in Pharmacology. Work from our laboratory that is cited in this paper has been supported by grant from DOD (PR130153) to S.M.N. We are thankful to the University of Toledo research programs as part of this work is funded by The University of Toledo's intramural startup fund for W.A.A.

REFERENCES

- [1] Wheatley DN. Landmarks in the first hundred years of primary (9+0) cilium research. *Cell Biol Int* 2005; 29(5): 333-9.
- [2] Nauli SM, Alenghat FJ, Luo Y, *et al.* Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet* 2003; 33(2): 129-37.
- [3] Cano DA, Sekine S, Hebrok M. Primary cilia deletion in pancreatic epithelial cells results in cyst formation and pancreatitis. *Gastroenterology* 2006; 131(6): 1856-69.
- [4] Masyuk AI, Masyuk TV, Splinter PL, Huang BQ, Stroope AJ, LaRusso NF. Cholangiocyte cilia detect changes in luminal fluid flow and transmit them into intracellular Ca²⁺ and cAMP signaling. *Gastroenterology* 2006; 131(3): 911-20.
- [5] Nauli SM, Kawanabe Y, Kaminski JJ, Pearce WJ, Ingber DE, Zhou J. Endothelial cilia are fluid shear sensors that regulate calcium signaling and nitric oxide production through polycystin-1. *Circulation* 2008; 117(9): 1161-71.
- [6] Van der Heiden K, Hierck BP, Krams R, *et al.* Endothelial primary cilia in areas of disturbed flow are at the base of atherosclerosis. *Atherosclerosis* 2008; 196(2): 542-50.
- [7] AbouAlaiwi WA, Takahashi M, Mell BR, *et al.* Ciliary polycystin-2 is a mechanosensitive calcium channel involved in nitric oxide signaling cascades. *Circ Res* 2009; 104(7): 860-9.
- [8] Nonaka S, Tanaka Y, Okada Y, *et al.* Randomization of left-right asymmetry due to loss of nodal cilia generating leftward flow of extraembryonic fluid in mice lacking KIF3B motor protein. *Cell* 1998; 95(6): 829-37.
- [9] Nonaka S, Shiratori H, Saijoh Y, Hamada H. Determination of left-right patterning of the mouse embryo by artificial nodal flow. *Nature* 2002; 418(6893): 96-9.

- [10] Essner JJ, Vogan KJ, Wagner MK, Tabin CJ, Yost HJ, Brueckner M. Conserved function for embryonic nodal cilia. *Nature* 2002; 418(6893): 37-8.
- [11] McGrath J, Somlo S, Makova S, Tian X, Brueckner M. Two populations of node monocilia initiate left-right asymmetry in the mouse. *Cell* 2003; 114(1): 61-73.
- [12] Karcher C, Fischer A, Schweickert A, *et al.* Lack of a laterality phenotype in Pkd1 knock-out embryos correlates with absence of polycystin-1 in nodal cilia. *Differentiation* 2005; 73(8): 425-32.
- [13] Andrade YN, Fernandes J, Vazquez E, *et al.* TRPV4 channel is involved in the coupling of fluid viscosity changes to epithelial ciliary activity. *J Cell Biol* 2005; 168(6): 869-74.
- [14] Teilmann SC, Byskov AG, Pedersen PA, Wheatley DN, Pazour GJ, Christensen ST. Localization of transient receptor potential ion channels in primary and motile cilia of the female murine reproductive organs. *Mol Reprod Dev* 2005; 71(4): 444-52.
- [15] Teilmann SC, Christensen ST. Localization of the angiotensin receptors Tie-1 and Tie-2 on the primary cilia in the female reproductive organs. *Cell Biol Int* 2005; 29(5): 340-6.
- [16] Hearn T, Spalluto C, Phillips VJ, *et al.* Subcellular localization of ALMS1 supports involvement of centrosome and basal body dysfunction in the pathogenesis of obesity, insulin resistance, and type 2 diabetes. *Diabetes* 2005; 54(5): 1581-7.
- [17] Winklbauer ME, Schafer JC, Haycraft CJ, Swoboda P, Yoder BK. The *C. elegans* homologs of nephrocystin-1 and nephrocystin-4 are cilia transition zone proteins involved in chemosensory perception. *J Cell Sci* 2005; 118(Pt 23): 5575-87.
- [18] Davenport JR, Watts AJ, Roper VC, *et al.* Disruption of intraflagellar transport in adult mice leads to obesity and slow-onset cystic kidney disease. *Curr Biol* 2007; 17(18): 1586-94.
- [19] Shah AS, Ben-Shahar Y, Moninger TO, Kline JN, Welsh MJ. Motile cilia of human airway epithelia are chemosensory. *Science* 2009; 325(5944): 1131-4.
- [20] Nishimura DY, Fath M, Mullins RF, *et al.* Bbs2-null mice have neurosensory deficits, a defect in social dominance, and retinopathy associated with mislocalization of rhodopsin. *Proc Natl Acad Sci U S A* 2004; 101(47): 16588-93.
- [21] Moore A, Escudier E, Roger G, *et al.* RPGR is mutated in patients with a complex X linked phenotype combining primary ciliary dyskinesia and retinitis pigmentosa. *J Med Genet* 2006; 43(4): 326-33.
- [22] Ghosh AK, Murga-Zamalloa CA, Chan L, Hitchcock PF, Swaroop A, Khanna H. Human retinopathy-associated ciliary protein retinitis pigmentosa GTPase regulator mediates cilia-dependent vertebrate development. *Hum Mol Genet* 2010; 19(1): 90-8.
- [23] Kulaga HM, Leitch CC, Eichers ER, *et al.* Loss of BBS proteins causes anosmia in humans and defects in olfactory cilia structure and function in the mouse. *Nat Genet* 2004; 36(9): 994-8.
- [24] Layman WS, McEwen DP, Beyer LA, *et al.* Defects in neural stem cell proliferation and olfaction in Chd7 deficient mice indicate a mechanism for hyposmia in human CHARGE syndrome. *Hum Mol Genet* 2009; 18(11): 1909-23.
- [25] Moorman SJ, Shorr AZ. The primary cilium as a gravitational force transducer and a regulator of transcriptional noise. *Dev Dyn* 2008; 237(8): 1955-9.
- [26] Christensen ST, Pedersen SF, Satir P, Veland IR, Schneider L. The primary cilium coordinates signaling pathways in cell cycle control and migration during development and tissue repair. *Curr Top Dev Biol* 2008; 85: 261-301.
- [27] Han YG, Kim HJ, Dlugosz AA, Ellison DW, Gilbertson RJ, Alvarez-Buylla A. Dual and opposing roles of primary cilia in medulloblastoma development. *Nat Med* 2009; 15(9): 1062-5.
- [28] Wong SY, Seol AD, So PL, *et al.* Primary cilia can both mediate and suppress Hedgehog pathway-dependent tumorigenesis. *Nat Med* 2009; 15(9): 1055-61.
- [29] Salathe M. Regulation of mammalian ciliary beating. *Annu Rev Physiol* 2007; 69: 401-22.
- [30] Zariwala MA, Knowles MR, Omran H. Genetic defects in ciliary structure and function. *Annu Rev Physiol* 2007; 69: 423-50.
- [31] Brunner S, Colman D, Travis AJ, *et al.* Overexpression of RPGR leads to male infertility in mice due to defects in flagellar assembly. *Biol Reprod* 2008; 79(4): 608-17.
- [32] Lee L, Campagna DR, Pinkus JL, *et al.* Primary ciliary dyskinesia in mice lacking the novel ciliary protein Pcdp1. *Mol Cell Biol* 2008; 28(3): 949-57.
- [33] Imai H, Hakkaku N, Iwamoto R, *et al.* Depletion of selenoprotein GPx4 in spermatocytes causes male infertility in mice. *J Biol Chem* 2009; 284(47): 32522-32.
- [34] Ibanez-Tallon I, Pagenstecher A, Fliegauf M, *et al.* Dysfunction of axonemal dynein heavy chain Mdnah5 inhibits ependymal flow and reveals a novel mechanism for hydrocephalus formation. *Hum Mol Genet* 2004; 13(18): 2133-41.
- [35] Sawamoto K, Wichterle H, Gonzalez-Perez O, *et al.* New neurons follow the flow of cerebrospinal fluid in the adult brain. *Science* 2006; 311(5761): 629-32.
- [36] Wodarczyk C, Rowe I, Chiaravalli M, Pema M, Qian F, Boletta A. A novel mouse model reveals that polycystin-1 deficiency in ependyma and choroid plexus results in dysfunctional cilia and hydrocephalus. *PLoS One* 2009; 4(9): e7137.
- [37] Eddy CA, Pauerstein CJ. Anatomy and physiology of the fallopian tube. *Clin Obstet Gynecol* 1980; 23(4): 1177-93.
- [38] Lyons RA, Saridogan E, Djahanbakhch O. The reproductive significance of human Fallopian tube cilia. *Hum Reprod Update* 2006; 12(4): 363-72.
- [39] Littlewood Evans A, Muller U. Stereocilia defects in the sensory hair cells of the inner ear in mice deficient in integrin alpha8beta1. *Nat Genet* 2000; 24(4): 424-8.
- [40] Grillet N, Kazmierczak P, Xiong W, *et al.* The mechanotransduction machinery of hair cells. *Sci Signal* 2009; 2(85): pt5.
- [41] Grillet N, Schwander M, Hildebrand MS, *et al.* Mutations in LOXHD1, an evolutionarily conserved stereociliary protein, disrupt hair cell function in mice and cause progressive hearing loss in humans. *Am J Hum Genet* 2009; 85(3): 328-37.
- [42] Nauli SM, Haymour HS, AbouAlaiwi WA, Lo ST, Nauli AM. Primary cilia are mechanosensory organelles in vestibular tissues. *Mechanosensitivity and Mechanotransduction*: Springer Netherlands; 2011. p. 317-50.
- [43] Collin SP, Barry Collin H. Primary cilia in vertebrate corneal endothelial cells. *Cell Biol Int* 2004; 28(2): 125-30.
- [44] Doughty MJ. Changes in cell surface primary cilia and microvilli concurrent with measurements of fluid flow across the rabbit corneal endothelium *ex vivo*. *Tissue Cell* 1998; 30(6): 634-43.
- [45] Kojimahara M. Endothelial cilia in rat mesenteric arteries and intramyocardial capillaries. *Z Mikrosk Anat Forsch* 1990; 104(3): 412-6.
- [46] Gallagher BC. Primary cilia of the corneal endothelium. *Am J Anat* 1980; 159(4): 475-84.
- [47] Satir P, Christensen ST. Overview of structure and function of mammalian cilia. *Annu Rev Physiol* 2007; 69: 377-400.
- [48] Hildebrandt F, Otto E. Cilia and centrosomes: a unifying pathogenic concept for cystic kidney disease? *Nat Rev Genet* 2005; 6(12): 928-40.
- [49] Praetorius HA, Spring KR. A physiological view of the primary cilium. *Annu Rev Physiol* 2005; 67: 515-29.
- [50] Berbari NF, Bishop GA, Askwith CC, Lewis JS, Mykytyn K. Hippocampal neurons possess primary cilia in culture. *J Neurosci Res* 2007; 85(5): 1095-100.
- [51] Whitfield JF. The neuronal primary cilium--an extrasynaptic signaling device. *Cell Signal* 2004; 16(7): 763-7.
- [52] Fuchs JL, Schwark HD. Neuronal primary cilia: a review. *Cell Biol Int* 2004; 28(2): 111-8.
- [53] Handel M, Schulz S, Stanarius A, *et al.* Selective targeting of somatostatin receptor 3 to neuronal cilia. *Neuroscience* 1999; 89(3): 909-26.
- [54] Malone AM, Anderson CT, Tummala P, *et al.* Primary cilia mediate mechanosensing in bone cells by a calcium-independent mechanism. *Proc Natl Acad Sci U S A* 2007; 104(33): 13325-30.

- [55] Whitfield JF. The solitary (primary) cilium--a mechanosensory toggle switch in bone and cartilage cells. *Cell Signal* 2008; 20(6): 1019-24.
- [56] Xiao Z, Zhang S, Magenheimer BS, Luo J, Quarles LD. Polycystin-1 regulates skeletogenesis through stimulation of the osteoblast-specific transcription factor RUNX2-II. *J Biol Chem* 2008; 283(18): 12624-34.
- [57] Christensen ST, Voss JW, Teilmann SC, Lambert IH. High expression of the taurine transporter TauT in primary cilia of NIH3T3 fibroblasts. *Cell Biol Int* 2005; 29(5): 347-51.
- [58] Schneider L, Clement CA, Teilmann SC, *et al.* PDGFRalpha signaling is regulated through the primary cilium in fibroblasts. *Curr Biol* 2005; 15(20): 1861-6.
- [59] Schroder JM, Schneider L, Christensen ST, Pedersen LB. EBI is required for primary cilia assembly in fibroblasts. *Curr Biol* 2007; 17(13): 1134-9.
- [60] Nauli SM, Haymour HS, Aboualaiwi WA, Lo ST, Nauli AM. Primary Cilia are Mechanosensory Organelles in Vestibular Tissues. In: A. Kamkin IK, editor. *Mechanosensitivity and Mechanotransduction*: Springer Science+Business Media; 2011. p. 317-50.
- [61] Schwartz EA, Leonard ML, Bizios R, Bowser SS. Analysis and modeling of the primary cilium bending response to fluid shear. *Am J Physiol* 1997; 272(1 Pt 2): F132-8.
- [62] Praetorius HA, Spring KR. Removal of the MDCK cell primary cilium abolishes flow sensing. *J Membr Biol* 2003; 191(1): 69-76.
- [63] Praetorius HA, Spring KR. Bending the MDCK cell primary cilium increases intracellular calcium. *J Membr Biol* 2001; 184(1): 71-9.
- [64] Kottgen M, Buchholz B, Garcia-Gonzalez MA, *et al.* TRPP2 and TRPV4 form a polymodal sensory channel complex. *J Cell Biol* 2008; 182(3): 437-47.
- [65] Wang S, Zhang J, Nauli SM, *et al.* Fibrocystin/polyductin, found in the same protein complex with polycystin-2, regulates calcium responses in kidney epithelia. *Mol Cell Biol* 2007; 27(8): 3241-52.
- [66] Hirokawa N, Tanaka Y, Okada Y, Takeda S. Nodal flow and the generation of left-right asymmetry. *Cell* 2006; 125(1): 33-45.
- [67] Jain R, Pan J, Driscoll JA, *et al.* Temporal relationship between primary and motile ciliogenesis in airway epithelial cells. *Am J Respir Cell Mol Biol* 2010; 43(6): 731-9.
- [68] Ally A. Ventrolateral medullary control of cardiovascular activity during muscle contraction. *Neurosci Biobehav Rev* 1998; 23(1): 65-86.
- [69] Krimer LS, Muly EC, 3rd, Williams GV, Goldman-Rakic PS. Dopaminergic regulation of cerebral cortical microcirculation. *Nat Neurosci* 1998; 1(4): 286-9.
- [70] Korner PI, Head GA, Badoer E, Bobik A, Angus JA. Role of brain amine transmitters and some neuromodulators in blood pressure, heart rate, and baroreflex control. *J Cardiovasc Pharmacol* 1987; 10(Suppl 12): S26-32.
- [71] Taylor EW, Jordan D, Coote JH. Central control of the cardiovascular and respiratory systems and their interactions in vertebrates. *Physiol Rev* 1999; 79(3): 855-916.
- [72] Greisen G. Autoregulation of cerebral blood flow in newborn babies. *Early Hum Dev* 2005; 81(5): 423-8.
- [73] Bolz S, Pieperhoff S, De Wit C, Pohl U. Chronic increases in transmural pressure reduce NO-mediated dilations in isolated resistance arteries of the hamster. *Acta Physiol Scand* 2000; 168(1): 113-7.
- [74] Rubanyi GM. Ionic mechanisms involved in the flow- and pressure-sensor function of the endothelium. *Z Kardiol* 1991; 80(Suppl 7): 91-4.
- [75] Spurrell BE, Murphy TV, Hill MA. Intraluminal pressure stimulates MAPK phosphorylation in arterioles: temporal dissociation from myogenic contractile response. *Am J Physiol Heart Circ Physiol* 2003; 285(4): H1764-73.
- [76] Pyke KE, Tschakovsky ME. The relationship between shear stress and flow-mediated dilatation: implications for the assessment of endothelial function. *J Physiol* 2005; 568(Pt 2): 357-69.
- [77] Raitakari OT, Celermajer DS. Flow-mediated dilatation. *Br J Clin Pharmacol* 2000; 50(5): 397-404.
- [78] Chien S. Mechanotransduction and endothelial cell homeostasis: the wisdom of the cell. *Am J Physiol Heart Circ Physiol* 2007; 292(3): H1209-24.
- [79] Brown TD. Techniques for mechanical stimulation of cells *in vitro*: a review. *J Biomech* 2000; 33(1): 3-14.
- [80] Stossel TP, Condeelis J, Cooley L, *et al.* Filamins as integrators of cell mechanics and signalling. *Nat Rev Mol Cell Biol* 2001; 2(2): 138-45.
- [81] Janmey PA, McCulloch CA. Cell mechanics: integrating cell responses to mechanical stimuli. *Annu Rev Biomed Eng* 2007; 9: 1-34.
- [82] Tarbell JM, Pahakis MY. Mechanotransduction and the glycocalyx. *J Intern Med* 2006; 259(4): 339-50.
- [83] Feletou M, Vanhoutte PM. Endothelium-dependent hyperpolarizations: past beliefs and present facts. *Ann Med* 2007; 39(7): 495-516.
- [84] Ignarro LJ, Cirino G, Casini A, Napoli C. Nitric oxide as a signaling molecule in the vascular system: an overview. *J Cardiovasc Pharmacol* 1999; 34(6): 879-86.
- [85] Nakane M. Soluble guanylyl cyclase: physiological role as an NO receptor and the potential molecular target for therapeutic application. *Clin Chem Lab Med* 2003; 41(7): 865-70.
- [86] Ong AC, Ward CJ, Butler RJ, *et al.* Coordinate expression of the autosomal dominant polycystic kidney disease proteins, polycystin-2 and polycystin-1, in normal and cystic tissue. *Am J Pathol* 1999; 154(6): 1721-9.
- [87] Hateboer N, Veldhuisen B, Peters D, *et al.* Location of mutations within the PKD2 gene influences clinical outcome. *Kidney Int* 2000; 57(4): 1444-51.
- [88] Prasad RM, Jin X, Aboualaiwi WA, Nauli SM. Real-time vascular mechanosensation through *ex vivo* artery perfusion. *Biol Proced Online* 2014; 16(1): 6.
- [89] Van der Heiden K, Groenendijk BC, Hierck BP, *et al.* Monocilia on chicken embryonic endocardium in low shear stress areas. *Dev Dyn* 2006; 235(1): 19-28.
- [90] Iomini C, Tejada K, Mo W, Vaananen H, Piperno G. Primary cilia of human endothelial cells disassemble under laminar shear stress. *J Cell Biol* 2004; 164(6): 811-7.
- [91] Poole CA, Jensen CG, Snyder JA, Gray CG, Hermanutz VL, Wheatley DN. Confocal analysis of primary cilia structure and colocalization with the Golgi apparatus in chondrocytes and aortic smooth muscle cells. *Cell Biol Int* 1997; 21(8): 483-94.
- [92] Wu J, Du H, Wang X, Mei C, Sieck GC, Qian Q. Characterization of primary cilia in human airway smooth muscle cells. *Chest* 2009; 136(2): 561-70.
- [93] Wu G, Markowitz GS, Li L, *et al.* Cardiac defects and renal failure in mice with targeted mutations in Pkd2. *Nat Genet* 2000; 24(1): 75-8.
- [94] Marszalek JR, Ruiz-Lozano P, Roberts E, Chien KR, Goldstein LS. Situs inversus and embryonic ciliary morphogenesis defects in mouse mutants lacking the KIF3A subunit of kinesin-II. *Proc Natl Acad Sci U S A* 1999; 96(9): 5043-8.
- [95] Sang L, Miller JJ, Corbit KC, *et al.* Mapping the NPHP-JBTS-MKS protein network reveals ciliopathy disease genes and pathways. *Cell* 2011; 145(4): 513-28.
- [96] Willaredt MA, Gorgas K, Gardner HA, Tucker KL. Multiple essential roles for primary cilia in heart development. *Cilia* 2012; 1(1): 23.
- [97] Clement CA, Ajbro KD, Koefoed K, *et al.* TGF-beta signaling is associated with endocytosis at the pocket region of the primary cilium. *Cell Rep* 2013; 3(6): 1806-14.
- [98] Kim K, Drummond I, Ibraghimov-Beskrovnya O, Klinger K, Arnaout MA. Polycystin 1 is required for the structural integrity of blood vessels. *Proc Natl Acad Sci U S A* 2000; 97(4): 1731-6.
- [99] Eder T, Schrier RW. Cardiovascular abnormalities in autosomal-dominant polycystic kidney disease. *Nat Rev Nephrol* 2009; 5(4): 221-8.
- [100] Griffin MD, Torres VE, Grande JP, Kumar R. Vascular expression of polycystin. *J Am Soc Nephrol* 1997; 8(4): 616-26.
- [101] Dolan JM, Meng H, Singh S, Paluch R, Kolega J. High fluid shear stress and spatial shear stress gradients affect endothelial

- proliferation, survival, and alignment. *Ann Biomed Eng* 2011; 39(6): 1620-31.
- [102] AbouAlaiwi WA, Muntean BS, Ratnam S, *et al.* Survivin-induced abnormal ploidy contributes to cystic kidney and aneurysm formation. *Circulation* 2014; 129(6): 660-72.
- [103] AbouAlaiwi WA, Ratnam S, Booth RL, Shah JV, Nauli SM. Endothelial cells from humans and mice with polycystic kidney disease are characterized by polyploidy and chromosome segregation defects through survivin down-regulation. *Hum Mol Genet* 2011; 20(2): 354-67.
- [104] Nazarewicz RR, Salazar G, Patrushev N, *et al.* Early endosomal antigen 1 (EEA1) is an obligate scaffold for angiotensin II-induced, PKC- α -dependent Akt activation in endosomes. *J Biol Chem* 2011; 286(4): 2886-95.
- [105] Li W, Wang H, Kuang CY, *et al.* An essential role for the Id1/PI3K/Akt/NF κ B/survivin signalling pathway in promoting the proliferation of endothelial progenitor cells *in vitro*. *Mol Cell Biochem* 2012; 363(1-2): 135-45.
- [106] Lin J, Guan Z, Wang C, *et al.* Inhibitor of differentiation 1 contributes to head and neck squamous cell carcinoma survival via the NF- κ B/survivin and phosphoinositide 3-kinase/Akt signaling pathways. *Clin Cancer Res* 2010; 16(1): 77-87.
- [107] Chapman AB, Rubinstein D, Hughes R, *et al.* Intracranial aneurysms in autosomal dominant polycystic kidney disease. *N Engl J Med* 1992; 327(13): 916-20.
- [108] Blitzer AL, Panagis L, Gusella GL, Danias J, Mlodzik M, Iomini C. Primary cilia dynamics instruct tissue patterning and repair of corneal endothelium. *Proc Natl Acad Sci U S A* 2011; 108(7): 2819-24.
- [109] Luo N, Conwell MD, Chen X, *et al.* Primary cilia signaling mediates intraocular pressure sensation. *Proc Natl Acad Sci U S A* 2014; 111(35): 12871-6.
- [110] Luo N, West CC, Murga-Zamalloa CA, *et al.* OCRL localizes to the primary cilium: a new role for cilia in Lowe syndrome. *Hum Mol Genet* 2012; 21(15): 3333-44.
- [111] Owens GK. Regulation of differentiation of vascular smooth muscle cells. *Physiol Rev* 1995; 75(3): 487-517.
- [112] Lu CJ, Du H, Wu J, *et al.* Non-random distribution and sensory functions of primary cilia in vascular smooth muscle cells. *Kidney Blood Press Res* 2008; 31(3): 171-84.
- [113] Shi ZD, Tarbell JM. Fluid flow mechanotransduction in vascular smooth muscle cells and fibroblasts. *Ann Biomed Eng* 2011; 39(6): 1608-19.
- [114] Fick GM, Johnson AM, Hammond WS, Gabow PA. Causes of death in autosomal dominant polycystic kidney disease. *J Am Soc Nephrol* 1995; 5(12): 2048-56.
- [115] Guay-Woodford LM. Renal cystic diseases: diverse phenotypes converge on the cilium/centrosome complex. *Pediatr Nephrol* 2006; 21(10): 1369-76.
- [116] Lilova MI, Petkov DL. Intracranial aneurysms in a child with autosomal recessive polycystic kidney disease. *Pediatr Nephrol* 2001; 16(12): 1030-2.
- [117] Biagini A, Maffei S, Baroni M, *et al.* Familial clustering of aortic dissection in polycystic kidney disease. *Am J Cardiol* 1993; 72(9): 741-2.
- [118] Capisonda R, Phan V, Traubuci J, Daneman A, Balfe JW, Guay-Woodford LM. Autosomal recessive polycystic kidney disease: outcomes from a single-center experience. *Pediatr Nephrol* 2003; 18(2): 119-26.
- [119] Gabow PA. Autosomal dominant polycystic kidney disease. *N Engl J Med* 1993; 329(5): 332-42.
- [120] Roy S, Dillon MJ, Trompeter RS, Barratt TM. Autosomal recessive polycystic kidney disease: long-term outcome of neonatal survivors. *Pediatr Nephrol* 1997; 11(3): 302-6.
- [121] Torres VE, Harris PC. Mechanisms of Disease: autosomal dominant and recessive polycystic kidney diseases. *Nat Clin Pract Nephrol* 2006; 2(1): 40-55; quiz.
- [122] Nauli SM, Rossetti S, Kolb RJ, *et al.* Loss of polycystin-1 in human cyst-lining epithelia leads to ciliary dysfunction. *J Am Soc Nephrol* 2006; 17(4): 1015-25.
- [123] AbouAlaiwi WA, Lo ST, Nauli SM. Primary Cilia: Highly Sophisticated Biological Sensors. *Sensors* 2009; 9(9): 7003-20.
- [124] Clausen P, Feldt-Rasmussen B, Iversen J, Lange M, Eidemak I, Strandgaard S. Flow-associated dilatatory capacity of the brachial artery is intact in early autosomal dominant polycystic kidney disease. *Am J Nephrol* 2006; 26(4): 335-9.
- [125] Wang D, Strandgaard S, Borresen ML, *et al.* Asymmetric dimethylarginine and lipid peroxidation products in early autosomal dominant polycystic kidney disease. *Am J Kidney Dis* 2008; 51(2): 184-91.
- [126] Gabow PA, Chapman AB, Johnson AM, *et al.* Renal structure and hypertension in autosomal dominant polycystic kidney disease. *Kidney Int* 1990; 38(6): 1177-80.
- [127] Gonzalo A, Gallego A, Rivera M, Orte L, Ortuno J. Influence of hypertension on early renal insufficiency in autosomal dominant polycystic kidney disease. *Nephron* 1996; 72(2): 225-30.
- [128] Chapman AB, Schrier RW. Pathogenesis of hypertension in autosomal dominant polycystic kidney disease. *Semin Nephrol* 1991; 11(6): 653-60.
- [129] Eder T, Schrier RW. Hypertension in autosomal-dominant polycystic kidney disease: early occurrence and unique aspects. *J Am Soc Nephrol* 2001; 12(1): 194-200.
- [130] Kolb RJ, Nauli SM. Ciliary dysfunction in polycystic kidney disease: an emerging model with polarizing potential. *Front Biosci* 2008; 13: 4451-66.
- [131] Abdul-Majeed S, Nauli SM. Calcium-mediated mechanisms of cystic expansion. *Biochim Biophys Acta* 2011; 1812(10): 1281-90.
- [132] Chapman AB, Johnson A, Gabow PA, Schrier RW. The renin-angiotensin-aldosterone system and autosomal dominant polycystic kidney disease. *N Engl J Med* 1990; 323(16): 1091-6.
- [133] Eder T, Chapman AB, Brosnahan GM, Edelstein CL, Johnson AM, Schrier RW. Effect of antihypertensive therapy on renal function and urinary albumin excretion in hypertensive patients with autosomal dominant polycystic kidney disease. *Am J Kidney Dis* 2000; 35(3): 427-32.
- [134] Ramunni A, Saracino A, Esposito T, Saliani MT, Coratelli P. Renal vascular resistance and renin-angiotensin system in the pathogenesis of early hypertension in autosomal dominant polycystic kidney disease. *Hypertens Res* 2004; 27(4): 221-5.
- [135] Kaspereit-Rittinghausen J, Deerberg F, Rapp KG, Weislo A. Renal hypertension in rats with hereditary polycystic kidney disease. *Z Versuchstierkd* 1990; 33(5): 201-4.
- [136] Lavoie JL, Lake-Bruse KD, Sigmund CD. Increased blood pressure in transgenic mice expressing both human renin and angiotensinogen in the renal proximal tubule. *Am J Physiol Renal Physiol* 2004; 286(5): F965-71.
- [137] Lohman-Adham M, Soto CE, Inagami T, Cassis L. The intrarenal renin-angiotensin system in autosomal dominant polycystic kidney disease. *Am J Physiol Renal Physiol* 2004; 287(4): F775-88.
- [138] Mancina G, Dell'Oro R, Quarti-Trevano F, Scopelliti F, Grassi G. Angiotensin-sympathetic system interactions in cardiovascular and metabolic disease. *J Hypertens Suppl* 2006; 24(1): S51-6.
- [139] van den Meiracker AH, Boomsma F. The angiotensin II-sympathetic nervous system connection. *J Hypertens* 2003; 21(8): 1453-4.
- [140] Chapman AB, Gabow PA. Hypertension in autosomal dominant polycystic kidney disease. *Kidney Int Suppl* 1997; 61: S71-3.
- [141] McPherson EA, Luo Z, Brown RA, *et al.* Chymase-like angiotensin II-generating activity in end-stage human autosomal dominant polycystic kidney disease. *J Am Soc Nephrol* 2004; 15(2): 493-500.
- [142] Shiroyanagi Y, Tanabe K, Hashimoto Y, *et al.* Kidney transplantation in the recipient with autosomal-dominant polycystic kidney disease: a single center experience. *Transplant Proc* 2000; 32(7): 1841-3.
- [143] Florijn KW, Chang PC, van der Woude FJ, van Bockel JH, van Saase JL. Long-term cardiovascular morbidity and mortality in autosomal dominant polycystic kidney disease patients after renal transplantation. *Transplantation* 1994; 57(1): 73-81.
- [144] Patch C, Charlton J, Roderick PJ, Gulliford MC. Use of antihypertensive medications and mortality of patients with autosomal dominant polycystic kidney disease: a population-based study. *Am J Kidney Dis* 2011; 57(6): 856-62.
- [145] Torres VE, Grantham JJ, Chapman AB, *et al.* Potentially modifiable factors affecting the progression of autosomal dominant

- polycystic kidney disease. Clin J Am Soc Nephrol 2011; 6(3): 640-7.
- [146] Torres VE, Bankir L, Grantham JJ. A case for water in the treatment of polycystic kidney disease. Clin J Am Soc Nephrol 2009; 4(6): 1140-50.
- [147] Chapman AB, Gabow PA, Schrier RW. Reversible renal failure associated with angiotensin-converting enzyme inhibitors in polycystic kidney disease. Ann Intern Med 1991; 115(10): 769-73.
- [148] Bursztyn M. Polycystic kidney disease and angiotensin-converting enzyme inhibitors. Ann Intern Med 1992; 117(1): 90.
- [149] Nutahara K, Higashihara E, Horie S, *et al.* Calcium channel blocker versus angiotensin II receptor blocker in autosomal dominant polycystic kidney disease. Nephron Clin Pract 2005; 99(1): c18-23.
- [150] Rahbari-Oskoui F, Williams O, Chapman A. Mechanisms and management of hypertension in autosomal dominant polycystic kidney disease. Nephrol Dial Transplant 2014; 29(12): 2194-201.
- [151] Schrier RW, Abebe KZ, Perrone RD, *et al.* Blood Pressure in Early Autosomal Dominant Polycystic Kidney Disease. N Engl J Med 2014; 371(24): 2255-66.
- [152] Abou Alaiwi WA, Lo ST, Nauli SM. Primary cilia: highly sophisticated biological sensors. Sensors (Basel) 2009; 9(9): 7003-20.
- [153] Nauli SM, Jin X, Hierck BP. The mechanosensory role of primary cilia in vascular hypertension. Int J Vasc Med 2011; 2011: 376281.


Graphical Abstract

Current Hypertension Reviews, 2015, Vol. 11, No. 1 00

Vascular Endothelial Primary Cilia: Mechanosensation and Hypertension

Ashraf M. Mohieldin, Hossain Saad Md Zubayer, Alzahra J. Al Omran, Hannah C. Saternos, Ali Zarban, Surya M. Nauli and Wissam A. AbouAlaiwi*

*Department of Pharmacology, University of Toledo Health Science Campus, Toledo, OH, USA



Primary cilia are mechanosensory organelles extending from the cell surface to sense cues in the extracellular environment and transduce signals into the cell producing a response that regulate a physiological phenomenon like blood pressure.